

# The Non-adherence with Treatment in Dialysis Patients in Iran, A Systematic Review

Ali Tayebi,<sup>1</sup> Behzad Einollahi,<sup>2</sup> Abolfazl Rahimi,<sup>3</sup> Masoud Sirati-Nir<sup>4</sup>

<sup>1</sup>Nephrology and Urology Research Center and Nursing Faculty, Baqiyatallah University of Medical Sciences, Tehran, Iran

<sup>2</sup>Nephrology and Urology Research Center, Baqiyatallah University of Medical Sciences, Tehran, Iran

<sup>3</sup>Nursing Faculty, Baqiyatallah University of Medical Sciences, Tehran, Iran

<sup>4</sup>Behavioral Sciences Research Center and Nursing Faculty, Baqiyatallah University of Medical Sciences, Tehran, Iran

**Keywords.** non-adherence, dialysis, patient, compliance

Non-adherence with treatment is a common problem among hemodialysis (HD) patients and was considered as a reliable factor in patients deteriorating, increasing admission chance and inappropriate responding to HD treatment. According to multiple factors related to Non-adherence to HD treatment and its importance in patients' quality of life, this study aimed at a systematic reviewing of Non-adherence with treatment in dialysis patients in Iran. Treatment with adherence status is one of the problems that patient has on dialysis. This study aims at detecting the most important causes of Non-adherence in patients on dialysis according to previous studies. First, all of the studies related to our title were searched using some keywords (dialysis, hemodialysis, adherence, and non-adherence) for English and Persian databases; Iranmedex, Magi ran, SID, Iran doc, Google, Google scholar, PubMed, Embase, CINAHL, PsycInfo, and Cochrane Database of Systematic that covering the period from 2010 to 2018 was reviewing following a predefined inclusion and exclusion criteria. As a result, 40 papers related to inclusion and exclusion criteria were identified and analyzed. Data were collected according to study features, measures of Non-adherence, prevalence rates and factors related to Non-adherence. The PRISMA (Preferred Reporting Items for Systematic Reviews and Meta-Analyses) guidelines were followed to conduct this systematic review. According the results of present study six main categories were found.

The main reasons for Non-adherence with treatment in dialysis patients listed in the papers included: Patient related factors, socioeconomic factors, Psychological factors, Health care related factors, Therapy related factors and Disease-related Factors below is provided for each explanation.

Treatment for improving adherence is so important and can play a very significant role to improve the health in patients on dialysis.

IJKD 2019;13:347-61  
www.ijkd.org

## INTRODUCTION

Non-adherence to treatment is a common problem among hemodialysis (HD) patients and was considered as a reliable factor in patients deteriorating, increasing admission chance and inappropriate responding to HD treatment. Kidney failure patients with low health literacy are at

increased risk in terms of the poor self-care and negative health outcome <sup>1</sup>. The high prevalence of kidney disease (KD) is considered as a main public health problem <sup>2</sup>.

In Iran, by the end of 2016, the population of patients with chronic renal failure who are treated by renal replacement therapy were about 58000,

among whom 29200 (94%) were being treated by HD and 1624 used peritoneal dialysis<sup>3</sup>. The reports express that, the rate of non-adherence with the treatment was different from 2 to 87%<sup>4</sup>. Non-adherence with the treatment is one of the most expensive processes which require an annual \$100billion costs<sup>5</sup>.

Non-adherence to extensive management of Chronic Kidney Disease (CKD) is a significant barrier to manage the population effectively. Interventions to improve adherence need to target the contributing factors to improve the quality of life<sup>6</sup>. Patients who are treated for Chronic Kidney Disease (CKD) are required to follow comprehensive treatment that is complicated and difficult to understand. Extensive management of CKD patients includes dialysis, medication, fluid and dietary restriction<sup>7</sup>.

Providing CKD extensive patient care in terms of dialysis plan, medication, fluid and dietary restriction is important to slow the progression and complications of CKD. However, evidence indicated that a substantial proportion of CKD patients were observed to deviate from prescribed dialytic, medication, dietary and fluid recommendations causing in progress challenges in the health care. Non-adherence rates to dialysis among CKD patients in previous studies were 2% to 87%<sup>8</sup>.

Patient adherence with the regimen is one of the behaviors related to chronic diseases, which predicts successful treatment and reduces the adverse effects and severity of the disease<sup>9</sup>. Adherence to dietary regimens is needed to the success of HD whose absence is related to significant health problems and complications for patients receiving HD. Besides educations, HD patients need a continuous care plan that improves their adherence knowledge, practice, and attitude<sup>10</sup>. Adherence to dietary regimens reduces the kidney workload and helps prevent renal complications and uremia<sup>11</sup>. Since adherence to prescribed therapies and prescriptions is a main challenge in patients with chronic diseases, if patients cannot comply with their treatment plans, these patients will suffer from severe consequences such as impaired development, resulting in the need for immediate treatment and hospitalization<sup>12</sup>. Non-adherence with the regimen in dialysis patients is related to frequent hospitalization, lack of benefits of treatment, high treatment costs and

a high number of physician visits<sup>13</sup>. Wabe *et al.* stated that adherence with the regimen is related to the decrease in glycosylated hemoglobin, which increases the rate of hemoglobin glycosylation by 14% to 16% with a 10% adherence with treatment<sup>14</sup>. The reason for Non-adherence is different in terms of the person's features and position. Many factors play a significant role in Non-adherence, and it is best to change the payment after identifying them with appropriate corrective actions<sup>15</sup>. A dynamic model, a patient adhered to treatment that has the necessary information, motivation and ability to control the disease, the relationship of information in Non-adherence, is a requirement to conduct the voluntary adherence behavior, is the patient's knowledge of treatment<sup>16,17</sup>. Therefore, this study was conducted with the aim of systematic review on the Non-adherence with Treatment in Dialysis Patients in Iran patients.

## MATERIALS AND METHODS

In this study, a systematic review of all researches conducted on determining the most important causes of Non-adherence in patients on dialysis was used according to previous studies till 2018.

### Literature Search

The method of expressing the data in this work includes determining the studied problem, collecting the data, analyzing and interpreting the findings based on the systematic study reporting system i.e. PRISMA. The above-mentioned protocol was used as a measure to search the papers. There was a time limit to perform the electronic searches (2010-2018). Some keywords (dialysis, hemodialysis, adherence, and non-adherence) searched for English and Persian databases; Iranmedex, Magiran, SID, Iran doc, Google, Google scholar, PubMed, Embase, CINAHL, PsycInfo, and Cochrane Database of Systematic Reviews to access the information requested from the studies related to our title.

In this study, the PRISMA guidelines were used for assessing the quality of Systematic Reviews.

### Inclusion and Exclusion Criteria

Inclusion and exclusion criteria were: papers related to adherence or Non-adherence, papers were in English and Persian, the papers were original and all the papers were free full text. As a result, 40 papers were found and analyses

related to inclusion and exclusion criteria. Data were collected based on studied characteristics, measures of Non-adherence, prevalence rates and factors related to Non-adherence. The PRISMA guidelines were followed in performing this systematic review.

**Data Extraction**


We excluded studies according to inclusion criteria such as papers related to adherence or Non-adherence, papers were in English and Persian, papers were original and all the papers were free full text that Figure shows the information.

**RESULTS**

According to the criteria to enter and leave, 40 papers were found. Thus, the main reasons for Non-adherence with treatment in dialysis patients listed in the papers included: Patient related factors, socioeconomic factors, Psychological factors, Health care related factors, Therapy related factors and Disease-related Factors below is provided for each explanation (Table 1). Table 2 summarizes the descriptive of included Studies.

**Table 1.** The main reasons for Non-adherence with treatment in Dialysis Patients

Categories of reasons contributing to Non-adherence	Sub reasons for each category
1. Patient related factors	Age Gender Level of education Health beliefs
2. Social support factors	Family and friends support Peer support <i>health system</i> support
3. Psychological factors	Depression Anxiety
4. Health care related factors	Availability of dialysis centers Appropriate educational system Attitudes of health workers Knowledge of health workers Facilities for dialysis centers
5. Therapy related factors	Muscle cramps Sleep disturbance Intra dialytic Hypotension Pain Physical fatigue
6. Disease-related Factors	Sexual dysfunction acute and chronic pains Uremia Osseous dysfunctions Sleeplessness



**Figure.** The selection process of studies included in this study.

**Table 2.** Descriptive Summary of Included Studies

No.	Author (year)	Study title	Country	Sample size	Study type	Results based on Main categories
1	Burns D. (2004)	Physical and psychosocial adaptation of blacks on hemodialysis	US	102	Non-experimental study	Putting trust in God was the most frequently identified strategy. The results from this study add to the body of knowledge about the problems that are experienced and the coping strategies in HD patients. <sup>67</sup>
2	Tsay SL. et al (2005)	Effects of an adaptation training programme for patients with end-stage renal disease	Taiwan	57	Randomized controlled trial	An adaptation training programme decreased stress and depression levels, and improved the quality of life of ATP patients receiving HD. <sup>68</sup>
3	Ekelund M-L. et al (2007)	Elucidating Issues Stressful for Patients in Predialysis and Dialysis: From Symptom to Context	Sweden	201	Descriptive-analytical study	The patients' behavior was avoidant or passive, although occasionally the account of being dependent upon others and the uncertainty associated with having to take special medication involved references to physical training as a way of coping. <sup>64</sup>
4	Herslman M. (2008)	Non-adherence to dietary prescriptions in chronic kidney disease	South Africa		Review of literature	The effect of obesity is thought to be related to its link with diabetes and hypertension, both known risk factors for chronic kidney disease. <sup>70</sup>
5	Rafii F. et al (2009)	Perceived social support in hemodialysis Patients	Iran	202	Cross-sectional, correlational	Perceptions in HD patients on social support and its related factors were varied. <sup>44</sup>
6	Sajjadi A. et al (2010)	Effective factors on fatigue in patients with chronic renal failure undergoing hemodialysis	Iran	56	Descriptive correlation study	By reducing the education level, income, increasing the age, dialysis history and risk of CRF the fatigue increased; These differences were significant only in terms of gender, income and history of disease. <sup>89</sup>
7	Bogner HR. et al (2012)	Integrated management of type 2 diabetes mellitus and depression treatment to improve medication adherence	Taiwan	31, 971	Randomized controlled trial	Depression is an important factor to reduce patient adherence. <sup>56</sup>
8	Scheurer D. et al (2012)	Association between different types of social support and medication adherence.	US	50 studies	Systematic review	This study found that practical social support was most consistently associated with greater medication adherence. <sup>88</sup>
9	Biniazi V. et al (2013)	Different Aspects of Fatigue Experienced by Patients Receiving Maintenance Dialysis in hemodialysis Units	Iran	163	Descriptive Cross-Sectional	Weakness and fatigue are among the debilitating complications of HD which impact on most patients who are continuously dialyzed. <sup>66</sup>
10	Vardanjani R. et al (2014)	Adherence to HD treatment and some related factors in HD patients	Iran	71	Descriptive analytic study	Treatment adherence condition in HD patients was moderate and related to chronic nature, economic problem and knowledge deficit. <sup>90</sup>
11	Montazeri RS, et al (2014)	Evaluation of Nutritional Knowledge in Terms of Dietary Sources of Protein, Phosphorous, Potassium and Fluids Restriction in hemodialysis Patients	Iran	50	Descriptive-analytical study	Patients' knowledge scores about dietary sources of phosphorous and potassium were significantly lower than those of other parts. <sup>91</sup>

Table 2. (Continued)

No.	Author (year)	Study title	Country	Sample size	Study type	Results based on Main categories
12	Farrokhi F. et al (2014)	Association between depression and mortality in patients receiving long-term dialysis	Iran	31 studies	Systematic review and meta-analysis	Association between depression and mortality and significant link was established between the depressive symptoms and mortality. <sup>58</sup>
13	Sayed M. et al (2014)	Assessment of socioeconomic burden of hemodialysis on ESRD patients in suez canal cities & ELArish	Egypt	121	Descriptive Cross-Sectional	The problems of transportation to and from the dialysis centers are other problems to improve the patient's conditions and better adherence with treatment. <sup>63</sup>
14	Joshi VD (2014)	Quality of life in end stage renal disease patients	India	109 studies	Systematic review	The end stage renal disease can be reduce the self-confidence and a sense of oneliness of the patient and increases the need to support. <sup>31</sup>
15	Al-Akhtari AM. et al(2014)	Medication adherence among adult patients on hemodialysis.	Saudi arabia	100	Descriptive Cross-Sectional	Gender, presence of a care-giver, number of members in the household and employment status have no impact on medication adherence. <sup>18</sup>
16	Ibrahim S. et al (2015)	Non-compliance among chronic hemodialysis patients and its impact on patients' outcomes	Egypt	100	Cross-sectional study	Non-adherence is associated with poor quality of life, depression and malnutrition. <sup>20</sup>
17	Qobadi M.et al (2015)	Health literacy and medical adherence in hemodialysis patients: the mediating role of disease-specific knowledge	Iran	204	Descriptive Cross-Sectional	There was a significant relation of kidney disease knowledge with both health literacy and adherence. <sup>1</sup>
18	Wim L. L. et al (2015)	Association of depressive and anxiety symptoms with adverse events in Dutch chronic kidney disease patients	Netherland	100	Prospective cohort study	Depressive symptoms were associated with an increased risk of poor clinical outcome. <sup>53</sup>
19	Munuo AE. et al (2016)	Nutrition knowledge, attitudes and practices among healthcare workers in management of chronic kidney diseases in selected hospitals	Tanzania	133	Cross-sectional study	Nutrition knowledge among participants was poor; though their attitude was positive they failed to practice due to poor knowledge. <sup>92</sup>
20	Janssen IM. Et al (2015)	Preferences of patients undergoing hemodialysis	Germany	4,518	Questionnaire-based study	Patients rate the absence of adverse effects of treatments, remaining in a good emotional state and good physical functioning. <sup>59</sup>
21	Loosman WL. et al (2015)	Iron supplementation associates with low mortality in pre-dialyzed advanced chronic kidney disease patients receiving erythropoiesis-stimulating agents	Taiwan	11,381	prospective cohort study	Depression increases the mortality and hospitalization in dialysis patients. <sup>54</sup>
22	Andrew S. et al(2016)	Educating Patients about CKD: The Path to Self-Management and Patient-Centered Care	USA	6 studies	Qualitative study	Considered the unawareness of the society and the lack of related educational centers and problems to access the scientific resources on renal failure, as the causes affecting the adherence with treatment. <sup>60</sup>
23	Rahimi Z. (2016)	Evaluation of Perceived Social Support of Patients Undergoing hemodialysis and Its Relation to Treatment Adherence and Clinical Outcomes in Educational Institutions in West Azerbaijan	Iran	255	Descriptive Cross-Sectional	There was relationship between perceived social support and adherence to diet restrictions, adherence to fluid restriction and adherence to medical prescriptions. <sup>93</sup>

Table 2. (Continued)

No.	Author (year)	Study title	Country	Sample size	Study type	Results based on Main categories
24	Wong MM et al (2017)	Interdialytic weight gain: trends, predictors, and associated outcomes in the international Dialysis Outcomes and Practice Patterns Study	12 countries	21,919	prospective cohort study	Gender impacted on the adherence with the treatment and males ignore to comply compared with females. <sup>22</sup>
25	Sinclair S.et al (2017)	Sympathy, empathy, and compassion: A grounded theory study of palliative care patients' understandings, experiences, and preferences	Canada	53	semi-structured interviews	There was relationship between the patient, the medical team and the family as one of the most effective factor on adherence with the treatment. <sup>29</sup>
26	Tannor EK et al (2017)	Quality of life in patients on chronic dialysis in South Africa	South Africa	106	comparative mixed methods study	Family support was impacted on the quality of life and adherence with treatment in chronic dialysis patients. <sup>38</sup>
27	Yusuke T. et al (2017)	Comparison of hospital mortality and readmission rates for Medicare patients treated by male vs female physicians	England	1,583, 028	Descriptive Cross-Sectional	Gender of the physicians is an effective factor on the mortality in patients, and female physicians treated the patients to have lower mortality. <sup>42</sup>
28	Tohme F. et al (2017)	Predictors and outcomes of non-adherence in patients receiving maintenance hemodialysis	US	286	(SMILE) trial	The novel factors independently associated with nonadherence to HD-related treatments, and independent associations of non-adherence with hospitalization and mortality. <sup>73</sup>
29	Kara B. (2018)	The Effect of Health Beliefs on the Observance of Diet and Refrain from Salt Consumption	Turkey	200	Cross-sectional study	Health beliefs related to salt restricted diet were independently associated with limiting salt intake, family support. <sup>28</sup>
30	Goh ZS, et al (2018)	Anxiety and depression in patients with end-stage renal disease	Singapore	Multiple studies	Narrative review	Social support as one of the emotional-coping mechanisms can impact on the quality of life. <sup>32</sup>
31	Ramezani T. et al (2018)	Effect of Educational Intervention Based on Self-efficacy Theory on Adherence to Dietary and Fluids-intake Restriction in HD Patients	Iran	70	Randomized clinical trial	The educational intervention based on Self-efficacy theory impacted on adherence to diet and fluid-intake restriction in patients with HD. <sup>94</sup>
32	Mousa I.et al (2018)	Dialysis-related factors affecting self efficacy and quality of life in patients on hemodialysis	Palestin	283	Descriptive Cross-Sectional	Impaired self-efficacy was related to the elderly, patients living with family, and patients with a high number of co-morbid diseases. <sup>36</sup>
33	Hedayati P.et al (2018)	Nonadherence behaviors and some related factors in kidney transplant recipients	Iran	183	Descriptive analytical study	The adherence to the treatment was associated with demographic characteristics and time passed since the transplant, levels of education, family support, marital status, and gender of the patients. <sup>39</sup>
34	Liu Y-M. et al (2018)	Role of resilience and social support in alleviating depression in patients receiving maintenance hemodialysis	Taiwan	194	Descriptive study	The severity of illness and daily living were the major determinants of depressive symptoms. High resilience could alleviate depressive symptoms in the older patients. <sup>40</sup>
35	Abdi A. et al (2018)	Prevalence of depression among Iranian patients under hemodialysis	Iran	24 studies	Systematic review and meta-analysis	More than half of the HD patients in Iran suffer from depression. Given the overlap of depression symptoms with uremia in HD patients. <sup>55</sup>

Table 2. (Continued)

No.	Author (year)	Study title	Country	Sample size	Study type	Results based on Main categories
36	Gerogianni G. et al (2018)	A holistic approach to factors affecting depression in hemodialysis patients	Greece	---	Review of literature	Depression is affected by non-adherence and co-morbid diseases. <sup>30</sup>
37	Janosevic D. et al (2018)	Difficult Patient Behavior in Dialysis Facilities	USA	---	Review of literature	realized that physicians the behavior and adherence of the patients could be managed through proper management and creation of appropriate environmental facilities. <sup>61</sup>
38	Pawar MAA. et al (2018)	Effectiveness of "music therapy" on "feeling of wellbeing" among patient undergoing hemodialysis	Indian	40	Cross sectional study	The positive effects of music on reducing anxiety and improving the physical and mental conditions of patients. <sup>62</sup>
39	Mukakarangwa MC. et al (2018)	Adherence to hemodialysis and Associated Factors among End Stage Renal Disease Patients	Rwanda	41	Descriptive Cross-Sectional	Age and religion. education of health care workers about the importance of not missing dialysis, perceived relative importance of HD, and experiencing difficulties during the procedure were associated with adherence to HD. <sup>74</sup>
40	Nagasawa H. et al (2018)	The Effect of Quality of Life on Medication Compliance Among Dialysis Patients	Japan	92	Descriptive Cross-Sectional	High sleep-related QOL is associated with better medication compliance. <sup>85</sup>

HD: hemodialysis

## DISCUSSION

Six categories of factors resulting in Non-adherence in CKD patients are patient related, psychological, disease related, socioeconomic, therapy related and health care system related factors. There are different sub factors for each category which results in Non-adherence as shown in Table 1.

### Patient Related Factors

Al-Akhtari (2014) considers age as one factor that affects the non-adherence with the treatment in HD patients. Despite that, in many studies, there has been only an insignificant effect of age on the adherence with the treatment<sup>18</sup>. However, in others, there was a positive and effective role of age on the adherence with the treatment<sup>19</sup>. Gender has also affected ineffectively the adherence with the treatment according to some studies<sup>20,21</sup>. In many studies, gender also affects effectively the adherence with the treatment and males mostly ignore to comply compared with females<sup>22, 23</sup>. Chironda (2016) expresses that, the level of education is one of the main factors affects the adherence with the treatment. Low level of education is related to the lack of knowledge about illness and treatment and

is a reason for non-adherence<sup>24</sup>. The knowledge of patients is a facilitator in adherence with the treatment<sup>25</sup>. The absence of health literacy is a factor to develop the health disorders and, thus, Non-adherence<sup>26,27</sup>. Belgüzar Kara (2018) conducted a cross sectional descriptive study entitled "The Effect of Health Beliefs on the Observance of Diet and Refrain from Salt Consumption" on 200 HD patients in Turkey. He concluded that a better understanding of the health beliefs about a limited-salt diet and the related factors may facilitate the implementation of appropriate interventions<sup>28</sup>.

Qobadi *et al.* (2015) aimed at examining the role of health literacy in patient's knowledge about kidney disease and medical care adherence and also to test the mediating role of kidney knowledge on the relationship between health literacy and medical adherence among patients treated with dialysis. This cross-sectional study was conducted on 240 consecutive patients diagnosed with end stage renal disease (ESRD) referred to dialysis ward affiliated to Tehran University of Medical Sciences from March 2014 to June 2014 according to the inclusion criteria. These findings show the high prevalence of low-health literacy in Iranian population and the health literacy is an important

point to promote the disease-specific knowledge and adherence behaviors among patients with dialysis<sup>1</sup>.

Sinclair *et al.* (2017) considered the relationship between the patient, the medical team and the family as one of the most effective factor on adherence with the treatment. Showing compassion and care by the treatment medical staff can be an important factor to encourage the patient to comply with the treatment<sup>29</sup>. Proper communication with and supporting the patient is one of the important adherence factors which is effective to control the depression in dialysis patients<sup>30</sup>.

### Social Support Factors

The duration of the treatment period and many problems encountered by these patients reduce their attention to family members and friends. However, the incidence of chronic renal failure and HD changes in one's life increases its dependency on others, reduces the self-confidence and a sense of loneliness of the patient and increases the need to support for others<sup>31</sup>. Social support as one of the emotional-coping mechanisms can affect the quality of life<sup>32</sup>. Social support is defined as a mental feeling about belonging, accepting, favoring and receiving assistance in the required conditions<sup>33</sup>. Understanding support is much more important than having it. In other words, the perception and attitude of patients towards received support is more important than the level of support provided to patients<sup>32</sup>. Identifying the benefits of social support can play an important role in the advanced stages of chronic renal failure, protecting the individual against stress-induced illness, reducing the psychological outcomes resulting from physical impairment, and improving compliance, tolerating chronic illness. Understanding social support can prevent the adverse physiological side effects of the disease, increase the level of self-care, affect positively the physical, psychological and social situation of the individual, and ultimately results in an increased performance<sup>34</sup>.

Information support also informed the person about how to care for themselves, control the disease and the diet, and this is an effective factor to improve the patient's condition, increase the self-confidence, and reduce his anxiety and fear<sup>35</sup>.

Mousa *et al* (2018) specify the factors related to the self-efficacy and health-related quality of life

(HRQoL) among HD patients. A cross-sectional descriptive correlation study was conducted on CKD patients treated with HD at 12 different dialysis centers in Palestine. The results show that impaired self-efficacy was related to the elderly, patients living with family, and patients with a high number of co-morbid diseases. Moreover, this study concluded that the worst HRQoL was related to patients with a low education level, lower levels of self-efficacy, and a high number of co-morbid diseases<sup>36</sup>.

Rahimi *et al.* (2017) stated that family support is one of the most important factors to increase the tolerance and resistance of patient and adherence with the treatment, which has been confirmed by various studies inside and outside Iran<sup>37</sup>. Elliott *et al.* also stated that family support is very effective on the quality of life and adherence with treatment in these patients due to the complicated and enormous problems of these patients<sup>38</sup>. Hedayati *et al.* believed that patients consider their partner as a key factor to support them. In fact, the support and preservation by the partner is the most important encouraging source for individuals during the illness<sup>39</sup>. The descriptive study of Leo *et al.* (2018) indicated that in HD patients, the spouse is the first source of support for the patient followed by children. They even reduced the severity of depression in these patients and finally result in the better adherence with the treatment<sup>40</sup>. Afkham *et al.* also stated that the physician is the first one in the medical team to confront with the patient and can help reduce the stress in the patient by having a supportive behavior at this stage. Participants in this study stated that the types of the support physicians can be mostly emotional (sympathy and giving hope) and informative<sup>41</sup>. Some studies have indicated the role of the gender of the physician and nurse to support the patients. In a descriptive study, Yusuke Tsugawa *et al.* realized that even the gender of the physicians is an effective factor on the mortality in patients, and female physicians treated the patients to have lower mortality. The researchers also expressed that the reason for this difference could be better communication with patients and better education by female physicians<sup>42</sup>. In their study, Aghaee *et al.* found that proper responsibility and training by nurses are the factors of hope and adherence with the treatment by the HD patients<sup>43</sup>. Perception of empathy provided by the nurses and



the needs of patients for sympathy and kindness affect the adherence with the treatment<sup>29</sup>. Rafi'ie *et al.* realized that friends' support is one of the important factors of adherence. These behaviors by friends play an important role to prevent the seclusion, isolation and depression of patients. This is due to this fact that HD patients cannot participate in group and collective activities due to changes in their physiognomy and the limitations to coordinate their activities with the members of the group. Therefore, they cut down their social relations and usually exclude from the group<sup>44</sup>.

Imani *et al.* (2013) also examined the effects of face-to-face education on body weight and some of the serum indices in patients treated with HD, and suggested heavily Nursing education was performed in dialysis department<sup>45</sup>. Hassanzadeh *et al.* (2011) also expressed that the effectiveness of two facial and video self-care educational programs on compliance with diet and fluids in HD patients was not significantly different<sup>46</sup>. Feizalahzadeh *et al* (2016) compare the effects of multimedia based education and traditional methods on life quality of HD patients. The HD and Essentials Care application based on the education can improve patients' life quality as well as traditional method. This application could be used to facilitate the patient education process to improve their life style<sup>47</sup>.

Al-Khattabi (2014) identifies the prevalence of adherence to (HD attendance, medications, fluid restrictions, and diet restrictions) among HD patients at governmental kidney centers<sup>48</sup>.

Montazeri and Sharifi (2014) conducted a study aims at measuring the nutritional knowledge in terms of dietary sources of protein, phosphorous, potassium and fluids restriction in HD patients. Patients' knowledge scores about dietary sources of phosphorous and potassium were significantly lower than those of other parts. It should be noted that qualified doctors, nurses and dieticians, should educate HD patients in relation to their literacy levels, especially when describing dietary sources of phosphorus, potassium and protein<sup>49</sup>.

Sa'adeh *et al.* (2018) measure the knowledge, attitudes and practices (KAP) of hypertensive patients towards prevention and early detection of CKD, and to determine the clinical and socio-demographic factors, which affect the KAP according to the prevention of CKD. Among

hypertensive patients, higher scores for total knowledge and attitudes in terms of prevention, male sex, and normal BMI were related to modestly higher scores to prevent the practices. Finally, the findings may encourage healthcare workers to consult better to improve knowledge<sup>50</sup>.

Adeline *et al* (2016) examined the nutrition knowledge, attitudes and current practices of healthcare workers to manage patients with CKD in selected hospitals. Nutrition knowledge among studied participants was poor; they failed to practice due to poor knowledge because their attitude was positive. Inadequate nutrition training in medical school, lack of resources and motivation were identified as factors that affect the nutrition management of CKD in the studied area. Recommendations provided by the study include: review of medical curriculum to incorporate clinical nutrition topics, continuous nutrition education programs for in-services, improving working conditions and hiring clinical nutritionists in each hospital department would improve nutrition management of CKD patients in hospitals<sup>51</sup>.

### Psychological Factors

Various studies have expressed that the main treatments for chronic renal failure, which include HD and kidney transplantation, both have psychological burden for the patient and can cause psychosocial problems for patients treated with these methods<sup>52</sup>.

In brief, depression and anxiety in renal patients are very common and depression in patients has several consequences such as lack of follow-up. It is estimated that depression in patients treated with dialysis is 25%<sup>53</sup>. Research expresses that depression increases the mortality and hospitalization in dialysis patients<sup>53, 54</sup>. Various factors such as age, sex, and duration of treatment can affect depression and anxiety in patients under 50 years of age. Also, alternative kidney therapies<sup>55</sup>.

Several studies have stated that hospitalized patients with depression also impose a higher cost to the health system. This can be due to the fact that depression is an important factor to reduce patient adherence<sup>56</sup>.

The absence of participation of depressed patients in treatment adds to their medical problems, endangers their health and in some cases even causes premature death<sup>57</sup>.

Farrokhi *et al.* (2014) systematically examine and analyze the relationship between depression and mortality risk in adults with kidney failure treated by long-term dialysis. There is significant relationship between-study heterogeneity in reports of depressive symptoms in dialysis patients, likely caused by high variability in the way to measure depressive symptoms. However, the total significant independent effect of depressive symptoms on survival of dialysis patients validates examining the fundamental mechanisms of this relationship and the possible benefits of interventions to improve depression on the outcomes<sup>58</sup>.

Abdi *et al* (2018) in a systematic review and meta-analysis examined the prevalence of depression in Iranian HD patients. The findings express that more than half of the HD patients in Iran suffer from depression. According to the overlap of depression symptoms with uremia in this group of patients, it is necessary to identify depression in these patients for early management and interventions<sup>55</sup>.

Janssen *et al* (2015) rated the relative importance of different outcomes for HD patients and to analyze whether the related importance is different among subgroups of patients. We assessed the relative importance of 23 outcomes as rated on a discrete visual analog scale within the framework. More important outcomes were hospital stays, accompanying symptoms, HD duration, and the improvement or preservation of a good emotional state. Outcomes related to the delivery or provision of care and aspects affecting the quality of life are rated by patients to be at least as important as clinical outcomes. Many of the outcomes assumed to be important by the patients are not regularly considered in research, evaluation studies, or quality programs<sup>59</sup>.

In a systematic study, Georgia Gerogianni *et al.* (2018) addressed the close relationship between depression and hospitalization, disability and mortality, job loss, suicidal thoughts, sleep disorders, fatigue, restless leg syndrome, sexual disorders, malnutrition and pain<sup>30</sup>.

### Health Care System Related Factors

The factors related to the health care system are another important factors affecting the adherence with treatment. Other factors that affect the adherence with treatment in this group of patients include the access to dialysis centers, problems and

costs of dialysis (drug procurement, transportation etc.), facilities and equipment of the dialysis centers and even the treatment of medical personnel. Andrew *et al.* (2016) considered the unawareness of the society and the lack of related educational centers and, of course, problems to access the scientific resources on renal failure, as the causes affecting the adherence with treatment. Another barrier is the absence of qualified educators in the community. Another important factor is the shortage of time among physicians and nurses to educate the patients in non-adherence. Another problem has been the lack of desire for education among patients. Another important barrier to adherence is also the disagreement on the starting time to educate the renal patients. System barriers to patient education are multi-faceted, and may have the absence of motivation for education, lack of support for decisions, and interdisciplinary health models. Another barrier is insufficient motivation and lack of willingness among the healthcare personnel<sup>60</sup>. In their research conducted on the facilities and behavior toward dialysis patients, Daniel *et al.* (2018) realized that physicians the behavior and adherence of the patients could be managed through proper management and other health personnel and creation of appropriate environmental facilities such as training, the emergence of the environment and proper behavior<sup>61</sup>. Pavar *et al.* (2018) examined the positive effects of music as a nursing intervention on reducing anxiety and improving the physical and mental conditions of patients<sup>62</sup>. Sayed *et al.* stated that the problems of transportation to and from the dialysis centers are other problems to improve the patient's conditions and better adherence with treatment<sup>63</sup>.

### Therapy-related Factors

Patients understand the complications of disease and HD as physical, social, and psychological problems. These problems will result in non-adherence with the treatment and they include: the physical complications of the illness, pain, fatigue and disability, food limitations, constant fear and anxiety, uncertainty, inadequacy in education, marriage and life, motherhood problems, and travel restrictions. In their study, Ekelund ML, Andersson (2007) examined the physical problems such as sleep disorders, complaints of cramps and itching in HD patients<sup>64</sup>. In Sajjadi *et al.* (2010),

most dialysis patients complained of high fatigue, and the mean of fatigue in the entire sample was 5.1 (out of the Likert score of 7)<sup>65</sup>. Biniazi *et al.* (2013) expressed that weakness and fatigue are among the debilitating complications of HD which affect most patients who are continuously dialyzed<sup>66</sup>. The pain of needles was another problem of HD that caused fear, anxiety and discomfort in patients. In his study, Burns D (2004) stated the need for frequent perforation of the vein and the related pain as one of the most important problems of HD patients<sup>67</sup>. Dietary limitation is another problem for many HD patients, which somewhat causes discomfort and harassment, and prevents the patient from having a normal life. Tsay *et al.* (2005) expressed the dietary constraints as one of the most stressing factors in HD patients<sup>68</sup>. Monahan *et al.* (2007) stated that the main social problems in a patient with chronic renal failure include social pressures, job loss, and restrictions on social and leisure activities<sup>69</sup>.

### Disease-related Factors

In HD patients, the complications of the illness are also related to the non-adherence. Herslman *et al.* explained the severe complications of Para clinical parameters such as anemia, calcium and phosphorus deficiency, and uremia and an increase in inflammatory chemical agents in these patients<sup>70</sup>. These mediators and complications reduce the tendency to eat and result in malnutrition<sup>71</sup>. Uremia causes frustration in the sense of taste and severe nutritional disorders<sup>72</sup>. Tehme *et al.* (2017) considered the acute and chronic pains as another cause of non-adherence in these patients. Sexual dysfunction is also a factor for non-adherence<sup>73</sup>. The incidence of various types of sexual disorders has been reported to be about 70% and 80%, which causes the stress in these patients<sup>74</sup>. Osseous dysfunctions, and especially an increase in the level of parathyroid hormone, are other factors that reduce the quality of life and consequently cause the non-adherence with the treatment<sup>74</sup>. The study conducted by Maung *et al.* (2017) showed that about 70 to 80% of the patients complained of sleeplessness. Sleeplessness leads to non-adherence with the treatment considering that sleeping is an effective factor in the ability of the body to control the stressful situations<sup>75</sup>.

Generally, Patients treated with HD are

encountering many challenges to adhere to treatment<sup>74</sup>. Beside all the factors that have been listed in various papers as causes of Non-adherence with the treatment of dialysis patients<sup>46, 76-79</sup>.

Depression is a type of mood disorder diagnosis by symptoms of low mood, decreased energy and interest, feeling guilty, difficulty in concentrating, anorexia, thoughts of death and suicide, insomnia or sleepiness, weight loss, and functional impairment<sup>80</sup>. Several studies have explained that hospitalized patients with depression also impose a higher cost to the health system<sup>81-84</sup>. This matter can be because of this fact that depression is a significant factor to reduce patient's adherence. The absence of participation of depressed patients in treatment adds to their medical problems, endangers their health and in some cases even causes premature death<sup>85</sup>. In contrast, the social support factor has been examined as the most important feature in the neutralization of undesirable effects of psychological pressures<sup>86</sup>. In addition to the positive effect that researchers have found on social support, it is also important to pay attention to understand the patient in this area. In other words, patients' perceptions and attitudes towards received support are more important than the patient's support<sup>87</sup>.

Social support is a social psychological factor that is identified as a facilitator of health behavior and can accelerate patient adherence with self-care behaviors<sup>88</sup>.

### CONCLUSION

In this study, we tried to examine the main causes of Non-adherence with the treatment of dialysis patients according to the studies; it is pointed out to factors such as psychosocial problems and especially depression, which play a significant role in isolating, confusing and disappointing individuals and as a consequence of Non-adherence with treatment. Other factors such as the lack of awareness and attitude of dialysis patients about their treatment and lack of adequate social support were also discussed.

Nurses, as experts who have the greatest relationship with patients during dialysis, can play a significant role by creating a continuous, dynamic care relationship to increase the awareness and practice of effective care for patients. On the other hand, economic problems, deprivation and difficult living conditions, side costs for patients

such as dialysis for dialysis, inability to work effectively, having diseases other than kidney problems and all are factors that cause many psychological problems in both patients and their families and it can cause many other problems, apart from enduring the pain and suffering of the disease. Therefore, government should designed the policies aim at minimizing the side effects of dialysis patients such as economic problems, cost of illness, etc.; in such a way that, the motivation and efforts of the patients are increased to continue life and improve the quality of life and follow-up of the treatment by reducing the side factors.

## REFERENCES

1. Qobadi M, Besharat MA, Rostami R, Rahiminezhad A. Health literacy and medical adherence in hemodialysis patients: the mediating role of disease-specific knowledge. *Thrita*. 2015;4(1).
2. Alikari V, Tsironi M, Matziou V, Babatsikou F, Psillakis KJIANHC. Adherence to Therapeutic Regimen in Adults Patients Undergoing Hemodialysis: The Role of Demographic and Clinical Characteristics. 2018;4:096.
3. Iran cd. A comprehensive report on the status of patients with renal failure in the world and Iran consortium dialysis Iran; 2017
4. Kim Y, Evangelista LS. Relationship between illness perceptions, treatment adherence, and clinical outcomes in patients on maintenance hemodialysis. *Nephrology nursing journal: journal of the American Nephrology Nurses' Association*. 2010;37(3):271.
5. Ghimire S, Castelino RL, Jose MD, Zaidi STR. Medication adherence perspectives in haemodialysis patients: a qualitative study. *BMC nephrology*. 2017;18(1):167.
6. Chironda G, Bhengu B. Contributing Factors to Non-Adherence among Chronic Kidney Disease (CKD) Patients: A Systematic Review of Literature. *Medical & Clinical Reviews*. 2016;2(04).
7. Short CD. Principles of Renal Replacement Therapy (RRT). *Imaging and Technology in Urology*: Springer; 2012: 339-342.
8. Griva K, Lai AY, Lim HA, Yu Z, Foo MWY, Newman SP. Non-adherence in patients on peritoneal dialysis: a systematic review. *PLoS One*. 2014;9(2):e89001.
9. Chironda G, Manwere A, Nyamakura R, Chipfuwa<sup>4</sup> T. Perceived health status and adherence to haemodialysis by End Stage Renal Disease patients: A case of a Central hospital in Zimbabwe.
10. Hashemi S, Tayebi A, Rahimi A, Einollahi B. Examining the effect of continuous care model on adherence to dietary regimen among patients receiving hemodialysis. *Iran J Crit Care Nurs*. 2015;7(4):215-220.
11. Mayers JD. Dietary restrictions in maintenance hemodialysis: experiences of English speaking West Indian adults. *Nephrology Nursing Journal*. 2000;27(3):315.
12. Deif HIA, ElSawi K, Selim M, NasrAllah MM. Effect of an Educational Program on Adherence to Therapeutic Regimen among Chronic Kidney Disease Stage5 (CKD5) Patients under Maintenance Hemodialysis. *Journal of Education and Practice*. 2015;6(5):21-33.
13. de Sa Basilio Lins SM, Leite JL, de Godoy S, Arimatea Branco Tavares JM, Rocha RG, Costa e Silva FV. Treatment adherence of chronic kidney disease patients on hemodialysis. *ACTA PAULISTA DE ENFERMAGEM*. 2018;31(1):54-60.
14. Wabe NT, Angamo MT, Hussein S. Medication adherence in diabetes mellitus and self management practices among type-2 diabetics in Ethiopia. *North American journal of medical sciences*. 2011;3(9):418.
15. Kim Y, Evangelista LS, Phillips LR, Pavlish C, Kopple JD. The End-Stage Renal Disease Adherence Questionnaire (ESRD-AQ): testing the psychometric properties in patients receiving in-center hemodialysis. *Nephrology nursing journal: journal of the American Nephrology Nurses' Association*. 2010;37(4):377.
16. Khalil AA, Darawad M, Al Gamal E, Hamdan-Mansour AM, Abed MA. Predictors of dietary and fluid non-adherence in Jordanian patients with end-stage renal disease receiving haemodialysis: a cross-sectional study. *Journal of Clinical Nursing*. 2013;22(1-2):127-136.
17. Som A, Groenendyk J, An T, et al. Improving dialysis adherence for high risk patients using automated messaging: Proof of concept. *Scientific Reports*. 2017;7(1):4177.
18. Alkatheri AM, Alyousif SM, Alshabanah N, et al. Medication adherence among adult patients on hemodialysis. *Saudi Journal of Kidney Diseases and Transplantation*. 2014;25(4):762.
19. Larsen J, Stovring H, Kragstrup J, Hansen DG. Can differences in medical drug compliance between European countries be explained by social factors: analyses based on data from the European Social Survey, round 2. *BMC Public Health*. 2009;9(1):145.
20. Ibrahim S, Hossam M, Belal D. Study of non-compliance among chronic hemodialysis patients and its impact on patients' outcomes. *Saudi Journal of Kidney Diseases and Transplantation*. 2015;26(2):243.
21. Bland RJ, Cottrell RR, Guylor LR. Medication compliance of hemodialysis patients and factors contributing to non-compliance. *Dialysis & Transplantation*. 2008;37(5):174-178.
22. Wong MM, McCullough KP, Bieber BA, et al. Interdialytic weight gain: trends, predictors, and associated outcomes in the international Dialysis Outcomes and Practice Patterns Study (DOPPS). *American Journal of Kidney Diseases*. 2017;69(3):367-379.
23. Takaki J, Nishi T, Shimoyama H, et al. Possible variances of blood urea nitrogen, serum potassium and phosphorus levels and interdialytic weight gain accounted for compliance of hemodialysis patients. *Journal of psychosomatic research*. 2003;55(6):525-529.
24. Chironda G, Bhengu B. Contributing factors to non-adherence among chronic kidney disease (CKD) patients: a systematic review of literature. *Medical & Clinical Reviews*. 2016;2(4):29.
25. Smith K, Coston M, Glock K, et al. Patient perspectives

- on fluid management in chronic hemodialysis. *Journal of Renal Nutrition*. 2010;20(5):334-341.
26. Lam LW, Twinn SF, Chan SW. Self-reported adherence to a therapeutic regimen among patients undergoing continuous ambulatory peritoneal dialysis. *Journal of advanced nursing*. 2010;66(4):763-773.
  27. Browne T, Merighi JR. Barriers to adult hemodialysis patients' self-management of oral medications. *American Journal of Kidney Diseases*. 2010;56(3):547-557.
  28. Kara B. Health beliefs related to salt-restricted diet and associated factors in Turkish patients on hemodialysis. *Journal of Transcultural Nursing*. 2018;29(2):155-164.
  29. Sinclair S, Beamer K, Hack TF, et al. Sympathy, empathy, and compassion: A grounded theory study of palliative care patients' understandings, experiences, and preferences. *Palliative medicine*. 2017;31(5):437-447.
  30. Gerogianni G, Kouzoupis A, Grapsa E. A holistic approach to factors affecting depression in haemodialysis patients. *International urology and nephrology*. 2018;50(8):1467-1476.
  31. Joshi VD. Quality of life in end stage renal disease patients. *World journal of nephrology*. 2014;3(4):308.
  32. Goh ZS, Griva K. Anxiety and depression in patients with end-stage renal disease: impact and management challenges—a narrative review. *International journal of nephrology and renovascular disease*. 2018;11:93.
  33. Shirazian S, Grant CD, Aina O, Mattana J, Khorassani F, Ricardo AC. Depression in chronic kidney disease and end-stage renal disease: Similarities and differences in diagnosis, epidemiology, and management. *Kidney International Reports*. 2017;2(1):94-107.
  34. Ladin K, Emerson J, Butt Z, et al. How important is social support in determining patients' suitability for transplantation? Results from a National Survey of Transplant Clinicians. *Journal of medical ethics*. 2018;44(10):666-674.
  35. Nixon AC, Bampouras TM, Pendleton N, Woywodt A, Mitra S, Dhaygude A. Frailty and chronic kidney disease: current evidence and continuing uncertainties. *Clinical kidney journal*. 2017;11(2):236-245.
  36. Mousa I, Ataba R, Al-ali K, Alkaiyat A, Sa'ed HZ. Dialysis-related factors affecting self-efficacy and quality of life in patients on haemodialysis: a cross-sectional study from Palestine. *Renal Replacement Therapy*. 2018;4(1):21.
  37. Z r. Evaluation of Perceived Social Support of Patients Undergoing Hemodialysis and Its Relation to Treatment Adherence and Clinical Outcomes in Educational Institutions in West Azerbaijan J Urmia Nurs Midwifery Fac. 2017;14(11):952-959.
  38. Tannor EK, Archer E, Kapembwa K, Van Schalkwyk SC, Davids MR. Quality of life in patients on chronic dialysis in South Africa: a comparative mixed methods study. *BMC nephrology*. 2017;18(1):4.
  39. Hedayati P, Shahgholian N, Ghadami A. Nonadherence behaviors and some related factors in kidney transplant recipients. *Iranian journal of nursing and midwifery research*. 2017;22(2):97.
  40. Liu Y-M, Chang H-J, Wang R-H, Yang L-K, Lu K-C, Hou Y-C. Role of resilience and social support in alleviating depression in patients receiving maintenance hemodialysis. *Therapeutics and clinical risk management*. 2018;14:441.
  41. Afkham Ebrahimi A, Nasr Esfahani M, Saghafi N. Patients' expectations and satisfaction with their treating physician. *Razi journal of medical sciences*. 2004;11(41):367-375.
  42. Tsugawa Y, Jena AB, Figueroa JF, Orav EJ, Blumenthal DM, Jha AK. Comparison of hospital mortality and readmission rates for Medicare patients treated by male vs female physicians. *JAMA internal medicine*. 2017;177(2):206-213.
  43. Aghei H. A look to nursing profession and their empathy effect with patients. *Keihan newspaper*. 2010;19623:5.
  44. Rafii F, Rambod M, Hosseini F. Perceived social support in hemodialysis Patients. *Journal of hayat*. 2009;15(1):5-12.
  45. Imani E, Khademi Z, Teshnizi SH, et al. The role of face-to-face education on body weight and some serum parameters in hemodialysis patients. 2013.
  46. Hadian Z, Rafiee Vardanjani L, Barimnejad L. The most important causes of non-adherence in patients on dialysis. *Clinical Excellence*. 2016;5(2).
  47. Feizalahzadeh H, Zagheri Tafreshi M, Moghaddasi H, Ashghali Farahani M, Zareh Z, Khalilzadeh M. Effectiveness of Multimedia Based on Education and Traditional Methods on Life Quality of Hemodialysis Patients. *Journal of Holistic Nursing And Midwifery*. 2016;26(2):69-78.
  48. Al-Khattabi GH. Prevalence of treatment adherence among attendance at hemodialysis in Makah. *Int J Med Sci Public Health*. 2014;3(5):592-598.
  49. Sharifi N, Montazeri RS. Evaluation of Nutritional Knowledge in Terms of Dietary Sources of Protein, Phosphorous, Potassium and Fluids Restriction in Hemodialysis Patients. *Jentashapir Journal of Health Research*. 2014;5(4):1-5.
  50. Sa'adeh HH, Darwazeh RN, Khalil AA, Sa'ed HZ. Knowledge, attitudes and practices of hypertensive patients towards prevention and early detection of chronic kidney disease: a cross sectional study from Palestine. *Clinical hypertension*. 2018;24(1):6.
  51. Munuo AE, Mugendi BW, Kisanga OA, Otieno GO. Nutrition knowledge, attitudes and practices among healthcare workers in management of chronic kidney diseases in selected hospitals in Dar es Salaam, Tanzania; a cross-sectional study. *BMC Nutrition*. 2016/01/16 2016;2(1).
  52. Stasiak CES, Bazan KS, Kuss RS, Schuinski AFM, Baroni G. Prevalence of anxiety and depression and its comorbidities in patients with chronic kidney disease on hemodialysis and peritoneal dialysis. *Jornal Brasileiro de Nefrologia*. 2014;36(3):325-331.
  53. Loosman WL, Rottier MA, Honig A, Siegert CE. Association of depressive and anxiety symptoms with adverse events in Dutch chronic kidney disease patients: a prospective cohort study. *BMC nephrology*. 2015;16(1):155.
  54. Kuo K-L, Hung S-C, Liu J-S, Chang Y-K, Hsu C-C, Tarng D-C. Iron supplementation associates with low mortality in pre-dialyzed advanced chronic kidney disease patients receiving erythropoiesis-stimulating agents: a nationwide

- database analysis. *Nephrology Dialysis Transplantation*. 2015;30(9):1518-1525.
55. Abdi A, Dalvand S, Vahedian-Azimi A, Ghanei Gheshlagh R, Kurdi A. Prevalence of depression among Iranian patients under hemodialysis; a systematic review and meta-analysis. *Journal of Nephrology*. 2018;7(4):224-232.
  56. Bogner HR, Morales KH, de Vries HF, Cappola AR. Integrated management of type 2 diabetes mellitus and depression treatment to improve medication adherence: a randomized controlled trial. *The Annals of Family Medicine*. 2012;10(1):15-22.
  57. Fernández A, Quan J, Moffet H, Parker MM, Schillinger D, Karter AJ. Adherence to newly prescribed diabetes medications among insured Latino and white patients with diabetes. *JAMA internal medicine*. 2017;177(3):371-379.
  58. Farrokhi F, Abedi N, Beyene J, Kurdyak P, Jassal SV. Association between depression and mortality in patients receiving long-term dialysis: a systematic review and meta-analysis. *American journal of kidney diseases*. 2014;63(4):623-635.
  59. Janssen IM, Gerhardus A, von Gersdorff GD, et al. Preferences of patients undergoing hemodialysis—results from a questionnaire-based study with 4,518 patients. Patient preference and adherence. 2015;9:847.
  60. Narva AS, Norton JM, Boulware LE. Educating patients about CKD: the path to self-management and patient-centered care. *Clinical Journal of the American Society of Nephrology*. 2016;11(4):694-703.
  61. Janosevic D, Wang AX, Wish JB. Difficult Patient Behavior in Dialysis Facilities. *Blood purification*. 2018:1-5.
  62. Pawar MAA, Jadhav MV, Patil MRV, Varghese MRA. EFFECTIVENESS OF" MUSIC THERAPY" ON" FEELING OF WELLBEING" AMONG PATIENT UNDERGOING HEMODIALYSIS IN TERTIARY CARE HOSPITAL, KARAD. *INDIAN JOURNAL OF APPLIED RESEARCH*. 2018;8(8).
  63. Sayed M, Abdelatty M, ELRawy B, et al. Assessment of socioeconomic burden of haemodialysis on ESRD patients in Suez canal cities and ELArish. *American Journal of Research Communication*. 2014;2(7):125-134.
  64. Ekelund M-L, Andersson SI. Elucidating issues stressful for patients in predialysis and dialysis: from symptom to context. *Journal of health psychology*. 2007;12(1):115-126.
  65. et.all As. Effective factors on fatigue in patients with chronic renal failure undergoing hemodialysis. *Critical care nursing*. 2014;3(1):33-38.
  66. Biniiaz V, Tayybi A, Nemati E, Shermeh MS, Ebadi A. Different aspects of fatigue experienced by patients receiving maintenance dialysis in hemodialysis units. *Nephro-urology monthly*. 2013;5(4):897.
  67. Burns D. Physical and psychosocial adaptation of blacks on hemodialysis. *Applied Nursing Research*. 2004;17(2):116-124.
  68. Tsay SL, Lee YC, Lee YC. Effects of an adaptation training programme for patients with end-stage renal disease. *Journal of Advanced Nursing*. 2005;50(1):39-46.
  69. Monahan F, Sands J, Neighbors M, Marek J, Green C. Phipps' medical-surgical nursing: health and illness perspectives. Philadelphia: St. Louis, MO: Mosby Elsevier. 2007.
  70. Herselman M. Non-adherence to dietary prescriptions in chronic kidney disease. *South African Journal of Clinical Nutrition*. 2008;21(2):13-14.
  71. Jha V, Garcia-Garcia G, Iseki K, et al. Chronic kidney disease: global dimension and perspectives. *The Lancet*. 2013;382(9888):260-272.
  72. Krenitsky J. Nutrition in renal failure: myths and management. *Practical gastroenterology*. 2004;28(9):40-59.
  73. Tohme F, Mor MK, Pena-Polanco J, et al. Predictors and outcomes of non-adherence in patients receiving maintenance hemodialysis. *International urology and nephrology*. 2017;49(8):1471-1479.
  74. Mukakarangwa MC, Chironda G, Bhengu B, Katende G. Adherence to Hemodialysis and Associated Factors among End Stage Renal Disease Patients at Selected Nephrology Units in Rwanda: A Descriptive Cross-Sectional Study. *Nursing research and practice*. 2018;2018.
  75. Maung S, Sara AE, Cohen D, Chapman C, Saggi S, Cukor D. Sleep disturbance and depressive affect in patients treated with haemodialysis. *Journal of renal care*. 2017;43(1):60-66.
  76. Verbrugghe M, Verhaeghe S, Lauwaert K, Beeckman D, Van Hecke A. Determinants and associated factors influencing medication adherence and persistence to oral anticancer drugs: a systematic review. *Cancer treatment reviews*. 2013;39(6):610-621.
  77. Wood L. A review on adherence management in patients on oral cancer therapies. *European Journal of Oncology Nursing*. 2012;16(4):432-438.
  78. Kondryn HJ, Edmondson CL, Hill J, Eden TO. Treatment non-adherence in teenage and young adult patients with cancer. *The lancet oncology*. 2011;12(1):100-108.
  79. Puts M, Tu H, Tourangeau A, et al. Factors influencing adherence to cancer treatment in older adults with cancer: a systematic review. *Annals of Oncology*. 2013;25(3):564-577.
  80. Sorait W. The association of self-efficacy and self-management behavior in adult patients with chronic kidney disease: an integrative review. *Journal of Kidney Treatment and Diagnosis*. 2018;1(1).
  81. Hamody ART, Kareem AK, Al-Yasri ARS, Ali AS. Depression in Iraqi hemodialysis patients. *Arab journal of nephrology and transplantation*. 2013;6(3):169-172.
  82. Markevičiūtė M, Matusevičiūtė A, Długauskas E, Danilevičiūtė V, Rimševičius L. Depresijos simptomų paplitimas tarp 3–5 stadijų lėtinę inkstų liga sergančių pacientų. *Medicinos teorija ir praktika*. 2016;22(3):228-232.
  83. King-Wing Ma T, Kam-Tao Li P. Depression in dialysis patients. *Nephrology*. 2016;21(8):639-646.
  84. Feroze U, Martin D, Reina-Patton A, Kalantar-Zadeh K, Kopple JD. Mental health, depression, and anxiety in patients on maintenance dialysis. *Iranian journal of kidney diseases*. 2010;4(3).
  85. Nagasawa H, Tachi T, Sugita I, et al. The Effect of Quality of Life on Medication Compliance Among Dialysis

- Patients. *Frontiers in Pharmacology*. 2018;9:488.
86. Mondesir FL, Carson AP, Durant RW, Lewis MW, Safford MM, Levitan EB. Association of functional and structural social support with medication adherence among individuals treated for coronary heart disease risk factors: Findings from the REasons for Geographic and Racial Differences in Stroke (REGARDS) study. *PloS one*. 2018;13(6):e0198578.
  87. Martos-Méndez MJ. Self-efficacy and adherence to treatment: the mediating effects of social support. *Journal of Behavior, Health & Social Issues*. 2015;7(2):19-29.
  88. Scheurer D, Choudhry N, Swanton KA, Matlin O, Shrank W. Association between different types of social support and medication adherence. *The American journal of managed care*. 2012;18(12):e461-467.
  89. Sajjadi A, Farmahini Farahani B, Esmailpoor Zanjani S, Dormanesh B, Zare M. Effective factors on fatigue in patients with chronic renal failure undergoing hemodialysis. *J Crit Care Nurs*. 2010;3(1):13-14.
  90. Vardanjani R, Parvin N, Mahmoodi Shan G, Molaie E, Hasheminia SAJJoCN, Midwifery. Adherence to hemodialysis treatment and some related factors in hemodialysis patients admitted in Shahrekord Hajar hospital. 2014;2(4).
  91. Montazeri RS, Sharifi N. Evaluation of nutritional knowledge in terms of dietary sources of protein, phosphorous, potassium and fluids restriction in hemodialysis patients. *Jentashapir Journal of Health Research*. 2014;5(4).
  92. Munuo AE, Mugendi BW, Kisanga OA, Otieno GO. Nutrition knowledge, attitudes and practices among healthcare workers in management of chronic kidney diseases in selected hospitals in Dar es Salaam, Tanzania; a cross-sectional study. *BMC Nutrition*. 2016;2(1):6.
  93. Rahimi ZJTJUNMF. Evaluation of Perceived Social Support of Patients Undergoing Hemodialysis and Its Relation to Treatment Adherence and Clinical Outcomes in Educational Institutions in West Azerbaijan Province, 2016. 2017;14(11):952-959.
  94. Ramezani T, Sharifirad G, Gharlipour Z, Mohebi S. Effect of educational intervention based on self-efficacy theory on adherence to dietary and fluids-intake restriction in hemodialysis patients. *Health Education and Health Promotion*. 2018;6(1):31-38.

Correspondence to:  
 Masoud Sirati-Nir  
 Behavioral Sciences Research Center and Nursing Faculty,  
 Baqiyatallah University of Medical Sciences, Tehran, Iran  
 E-mail: masoudsirati@gmail.com

Received December 2018  
 Revised April 2019  
 Accepted July 2019