### Accepted Manuscript

Liquid phase extraction of nanosized biologically active estrogenic pollutants by using an efficient adsorbent

Roohollah Hafizi, Ramezan Ali Taheri, Hamid Moghimi

PII:	S0167-7322(18)31425-9
DOI:	doi:10.1016/j.molliq.2018.05.079
Reference:	MOLLIQ 9134
To appear in:	Journal of Molecular Liquids
Received date:	17 March 2018
Revised date:	24 April 2018
Accepted date:	18 May 2018

Please cite this article as: Roohollah Hafizi, Ramezan Ali Taheri, Hamid Moghimi, Liquid phase extraction of nanosized biologically active estrogenic pollutants by using an efficient adsorbent. Molliq (2018), doi:10.1016/j.molliq.2018.05.079

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



# ACCEPTED MANUSCRIPT



Available online at www.sciencedirect.com



MOLLIQ

Journal of Molecular Liquids ? (2018) 1–14

# Liquid phase extraction of nanosized biologically active estrogenic pollutants by using an efficient adsorbent

Roohollah Hafizi<sup>a,b</sup>, Ramezan Ali Taheri<sup>b,\*</sup>, Hamid Moghimi<sup>c</sup>

<sup>a</sup>Department of Physics, Isfahan University of Technology, Isfahan, 84156-83111, Iran <sup>b</sup>Nanobiotechnology Research Center, Baqiyatallah University of Medical Sciences, Tehran, Iran <sup>c</sup>Department of Microbial Biotechnology, School of Biology, College of Science, University of Tehran, Tehran, Iran

#### Abstract

Dangerous nanosized micropollutants and estrogenic hormones which might severely affect the environment are more found in water effluents, recently, and detection of them is of high importance. As the result, reliable and inexpensive absorptive extraction and removal applications are favored. As the second most abundant biomolecule on earth, a lignocellulosic material can be a potential choice as nanosized adsorbent. In this paper, quantum chemistry techniques are employed as the state of the art approach for predictive modeling, to investigate the feasibility of using this material in extraction and removal of of eight estrogens including estradiol, estrone, testosterone, progesterone, estriol, mestranol, ethinylestradiol and diethylstilbestrol. Due to nano-scaled dimensions of these pollutants, removing them by using current commercial methods is challenging and a new approach needed to be developed. The mechanism of removing micropollutants from water affluent is based on adsorption. This fact motivated us to simulate the adsorption process, and check how well the adsorbent interact with nanosized micropollutants. In order to evaluate the affinity of lowest energy sites on structure of the lignocellulosic material toward each estrogenic hormones and micropollutants, we carried out configurational bias Monte Carlo search of the configurational space of the lignocellulosic material-estrogenic micropollutants systems according to a simulated annealing schedule. To check the response and sensitivity of the lignocellulosic material to each estrogenic micropollutants, a combination of modified Flory-Huggins model and molecular simulation techniques was used. Evaluation of detachment energy revealed the potential of the lignocellulosic material as a reusable material for removing these compounds in an aqueous media.

Keywords: estrogenic micropollutants, water effluent, biosensors, computational chemistry, molecular modeling

#### 1. Introduction

As a consequence of usage of hormonal drugs in disease treatments, various types of biologically active nanosized micropollutants such as estrogens are released into environment,

<sup>\*</sup>Corresponding Author

Email address: taheri@bmsu.ac.ir (Ramezan Ali Taheri)

https://daneshyari.com/article/7841903