

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/233942320>

Can atrial natriuretic peptides measurement diagnose heart failure at different age groups?

Article in *Journal of research in medical sciences* · December 2012

Source: PubMed

CITATIONS

0

READS

29

*Letter to Editor***Can atrial natriuretic peptides measurement diagnose heart failure at different age groups?***J Res Med Sci 2012; 17(1): 116-117*

With interest, I read the article by Khalilian et al.¹ entitled "Relationship between echocardiographic findings and laboratory serum biomarkers in patients with and without low cardiac output" that recently published in your valuable journal. We know that in adult patients, atrial natriuretic peptides (ANP) and brainnatriuretic peptides (BNP) have been used for diagnosis, monitoring of treatment effects, and estimating the prognosis of systolic and diastolic heart failure.² Furthermore, ANP would remain quite stable during storage in plasma at -80°C for even 12 months.³ I would like to attract the authors' and readers' attention to some points that can be helpful in conducting future studies.

First, it is known that patient characteristics, such as age, gender, weight, and glomerular filtration rate (GFR) may influence the ANP and BNP plasma levels.⁴ Therefore, it was better if the case and control groups were matched in gender, just similar to what was performed for age. In addition, the authors should have selected end-stage renal disease (ESRD) patients with normal systolic heart

function as the control group for comparing with the case group.

In this study, the highest BNP and ANP plasma concentrations were found in children and adolescence with systolic heart failure. However, elevation of BNP and ANP plasma levels were higher in patients with significant volume overload than those with isolated pressure overload. These findings were consistent with the results reported by Westerlind et al.⁵

Also, I would like to get your attention to an interesting point. Westerlind et al. showed a linear correlation between ANP plasma concentration and left ventricular ejection fraction (LVEF) in children and adolescents and Zolty et al. reported such correlation in adult patients (aged 31-62 years).³ But surprisingly, Rutten et al. reported that the BNP, and not the ANP plasma level increased progressively with the LVEF decline in geriatric outpatients (older than 65 years).⁴ This suggests that measurement of ANP plasma level in these patients may have no additional value compared with measurement of the BNP level. Thus, further studies are required for evaluation of this difference among the age groups.

*Allahyar Golabchi¹***Conflict of Interests**

Authors have no conflict of interests.

References

1. Khalilian MR, Sabri MR, Nikyar A, Haghjooy Javanmard Sh. Relationship between echocardiographic findings and laboratory serum biomarkers in patients with and without low cardiac output. *J Res Med Sci* 2011; 16(11): 1397-404.

1- Cardiologist, Health Research Center, Bagiyatallah University of Medical Sciences, Tehran, Iran.
E-mail: golabchi@edc.mui.ac.ir

2. Richards M, Nicholls MG, Espiner EA, Lainchbury JG, Troughton RW, Elliott J, et al. Comparison of B-type natriuretic peptides for assessment of cardiac function and prognosis in stable ischemic heart disease. *J Am Coll Cardiol* 2006; 47(1): 52-60.
3. Zolty R, Bauer C, Allen P, Garrity M, Vittorio TJ. Atrial natriuretic peptide stability. *Clin Biochem* 2008; 41(14-15): 1255-8.
4. Rutten JH, van d, V, van der Cammen TJ, ten Cate FJ, Vletter WB, Boomsma F, et al. Associations between plasma natriuretic peptides and echocardiographic abnormalities in geriatric outpatients. *Arch Gerontol Geriatr* 2008; 47(2): 189-99.
5. Westerlind A, Wahlander H, Berggren H, Lundberg PA, Holmgren D. Plasma levels of natriuretic peptide type B and A in children with heart disease with different types of cardiac load or systolic dysfunction. *Clin Physiol Funct Imaging* 2008; 28(4): 277-84.