

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/233955064>

# Survey the effect of aerobic exercise on aerobic capacity in patients with coronary artery disease (CAD)

Article in *Pakistan Journal of Medical Sciences Online* · October 2007

CITATIONS

6

READS

121

4 authors, including:



**Majid Najafi Kalyani**

Shiraz University of Medical Sciences

35 PUBLICATIONS 308 CITATIONS

SEE PROFILE



**Abbas Ebadi**

Baqiyatallah University of Medical Sciences

544 PUBLICATIONS 2,822 CITATIONS

SEE PROFILE



**Soheilnajafi Mehri**

Baqiyatallah University of Medical Sciences

25 PUBLICATIONS 180 CITATIONS

SEE PROFILE

Some of the authors of this publication are also working on these related projects:



Model of stress of exposure to sharps injury in Iranian nurses: A Path Analysis [View project](#)



Effects of Stress on Mothers of Hospitalized Children in a Hospital in Iran [View project](#)

## SURVEY THE EFFECT OF AEROBIC EXERCISE ON AEROBIC CAPACITY IN PATIENTS WITH CORONARY ARTERY DISEASE (CAD)

Majid Najafi Kalyani<sup>1</sup>, Abbas Ebad<sup>2</sup>,  
Soheil Najafi Mehri<sup>3</sup>, Mohammad Hasan Kalantar Motamedi<sup>4</sup>

### ABSTRACT

**Objective:** Increased aerobic exercise capacity appears to reduce both all-cause mortality and cardiovascular disease mortality. Physical exercise to improve maximal oxygen consumption ( $VO_{2max}$ ) is thus strongly recommended, however evidence regarding the most efficient training intensity for patients with coronary artery disease (CAD) is still lacking. The purpose of this randomized study was to assess the effects of aerobic exercise for increasing  $VO_{2max}$  in stable CAD-patients.

**Methodology:** Thirty stable CAD-patients were randomized to supervised walking 30 min three times a week for 10 weeks. Before and after training  $VO_{2max}$  was predicted from Bruce treadmill test.

**Results:** Before training  $VO_{2max}$  was  $35.2 \pm 4.32$  ml/kg/min and after training the mean  $VO_{2max}$  was  $43.1 \pm 3.4$  ml/kg/min. This difference was significant ( $p < 0.05$ ).

**Conclusions:** Aerobic exercise is effective for increasing  $VO_{2max}$  in stable CAD-patients. As  $VO_{2max}$  seems to reflect a continuum between health and cardiovascular disease and death, the present data may be useful in designing effective training programmes for improved health in the future.

**KEY WORDS:** Coronary artery disease, Aerobic exercise, Maximal oxygen consumption ( $VO_{2max}$ ).

Pak J Med Sci October - December 2007 (Part-I) Vol. 23 No. 5 665-670

1. Majid Najafi Kalyani, Instructor, MSN.  
Department of Medical Surgical,  
Faculty of Nursing,  
Baqiatallah Medical Science University,  
Tehran - Iran.
  2. Abbas Ebad, Instructor, MSN., Ph.D Candidate  
Department of Medical Surgical,  
Faculty of Nursing,  
Tarbiat Modares University,  
Baqiatallah Medical Science University  
Tehran - Iran.
  3. Soheil Najafi Mehri, Instructor, MSN., Ph.D Candidate,  
Department of Medical Surgical, Faculty of Nursing.
  4. Mohammad Hasan Kalantar Motamedi, M.D. & Cardiac  
Surgeon, Associated Professor
- 1,3-4: Baqiatallah Medical Science University,  
Tehran - Iran.

### Correspondence

Majid Najafi Kalyanim, MSN  
E-mail: majidnajafi5@yahoo.com

- \* Received for Publication: January 27, 2007
- \* Revision Received: January 31, 2007
- \* Revision Accepted: July 13, 2007

## INTRODUCTION

Higher levels of physical activity and fitness appear to reduce all-cause mortality and cardiovascular disease (CVD) mortality.<sup>1-6</sup> Physical exercise is thus strongly recommended in both primary and secondary prevention of CVD.<sup>7-11</sup> Peak aerobic exercise capacity is found to be the strongest independent predictor of mortality compared with other established risk factors among both healthy individuals and those with CVD<sup>12</sup> More specific, peak aerobic exercise capacity directly measured as maximal oxygen consumption ( $VO_{2max}$ ) was recently found to be the single best predictor of both cardiac and all-cause deaths among patients with established CVD<sup>13</sup>

For health promotions, patients with coronary artery disease (CAD) are recommended to regularly exercise at a mild to moderate

levels.<sup>7-9</sup> However, aerobic exercise training programmes are most often carried out at low-to-moderate intensities.<sup>14</sup> Both walking and vigorous exercise were found to be equally effective in increasing aerobic capacity and reduce cardiovascular risks.<sup>15-21</sup> However, two large cohort studies found that higher intensity of physical activity was related to reduced risk, as reflected by an inverse association between exercise intensity and coronary heart disease incidence in men.<sup>22,23</sup>

As aerobic exercise capacity seems to reflect a continuum between health and cardiovascular disease and death, it is important to design effective programmes for exercise-induced gains of  $VO_{2max}$  for patients with cardiovascular risk. Previous studies investigating the influence of different exercise intensities for improvements of  $VO_{2max}$ .<sup>14</sup> The aim of the present study was therefore to assess the effects aerobic exercise programmes of walking. The hypothesis was that aerobic exercise at low intensity (walking 30min three times a week for 10 weeks) is effective for increasing aerobic capacity in stable CAD-patients.

## METHODS

Thirty eligible participants were enrolled in the study in August 2005 and the study was accomplished in December 2006. All patients had undergone a medical investigation for CAD at Jamaran Heart Hospital of Baqiatallah Medical Science University, within one year prior to the study (Table-I). Inclusion criteria were angiographically documented CAD in at least one major epicardial vessel. In addition, subjects had clinical evidence of CAD in the form of previous myocardial infarction, significant stenosis treated with coronary artery bypass surgery (CABG) or percutaneous coronary intervention (PCI), or ischaemia in exercise-electrocardiogram (ECG).

The study was accomplished according to the Declaration of Helsinki. The regional committee for medical research ethics approved the study protocol. Written and informed consent was obtained from all subjects at the beginning

of the study. Before prediction of  $VO_{2max}$  the subjects were informed about the test, and instructed to exercise to their maximum limit. A standard 12-lead ECG was recorded at rest and at the end of each work level, and patients were stopped if any indication for terminating testing according to current guidelines took place.<sup>9</sup> To familiarize with Bruce treadmill test, the test started on a flat treadmill where participants learned to walk without grasping the handrails. As soon as they could walk properly, the speed and inclination was individually adjusted (2.74 -12.07km.h<sup>-1</sup> and 10–28%) for a 3-minute warm-up. After the warm-up period using a Bruce protocol where the speed and the incline was increased every three minute until HR>220-age or patient be fatigued. Patients were walking during the study. Heart rate was continuously recorded using a Polar Sport Tester (Polar Electro OY, Finland) and maximum attainable heart rate (HR peak) was determined.

The patients met for training three times per week for 10 weeks under supervision of an exercise physiologist. The patients were instructed not to add any leisure exercise during the study period. Training consisted of walking.

All values are expressed as mean±standard deviation (SD). Changes before and after training were assessed using the paired sample t test. A two-tailed P<0.05 was accepted as statistically significant for all tests.

## RESULTS

Patients demographics are given in Table-I with mean age of 45.2±5.6 years. The  $VO_{2max}$  increased significantly after training (Fig-1). Before training  $VO_{2max}$  was 35.2±4.32ml/kg/min and after training reached to 43.1±3.4ml/kg/min (Table-II). This difference was signifi-

Table-I: Patients demographics

Variable	Meanmean±SD	Max	Min
Age (Years)	45.2±5.6	32	54
Weight (kg)	72.3±6.2	62	83
Height (cm)	166.1±8.8	154	182
BMI	25.3±6.2	19.3	33.4

Table-II: Homodynamic parameters in patient with CAD

Status variable	Before training mean±SD	After training mean±SD
Vo <sub>2</sub> max ml/kg/min	35.2±4.32	43.1±3.4
Time Minutes	8.2±1.4	11.1±2.3
Distance Meters	822.3±120.3	1126.7±230.2

cant ( $p < 0.05$ ). The improvement after training was significantly greater compared to the before training. HR<sub>max</sub> after training significantly reduced. There were detectable changes in resting blood pressure and resting heart rate in patients after the training period. The average time for completing test before training was  $8.2 \pm 1.4$  minute and after training was  $11.1 \pm 2.3$  minutes (Fig-2). Total distance before training was  $822.3 \pm 120.3$  m and after training reached to  $1126.7 \pm 230.2$  m. This differences were significant ( $P < 0.05$ ). There were no episodes of cardiac events during the study.

## DISCUSSION

The results of this randomized controlled study demonstrate that aerobic exercise is superior for increasing  $VO_{2max}$  in stable CAD-patients. Although  $VO_{2max}$  increased after 10 weeks of training, the improvement was significantly larger in the men. With regard to the total amount of work performed, this study solely points out aerobic exercise as a key factor for increasing aerobic capacity in this




Figure-1: Maximal oxygen consumption ( $VO_{2max}$ ) before and after aerobic exercise training in patient with CAD.




Figure-2: Time before and after aerobic exercise training in patient with CAD

patient group. In view of the prognostic importance of increasing  $VO_{2max}$  for this patient group, aerobic exercise may be considered in future rehabilitation programmes.

The present study is one of few where CAD-patients are performing aerobic interval exercise 30 minutes three times a week for 10 weeks throughout the whole training period. Since all subjects exercised with a heart rate monitoring device, the load of the treadmill could progressively be adjusted to keep the relative exercise intensity constant as training adaptations occurred. The improvement of after training compared to before training reflects the importance of aerobic exercise when determining the increase of  $VO_{2max}$ . Two earlier studies involving CAD-patients have employed aerobic interval exercise with elements of the high intensity as in the present study, both with a tremendous increase of  $VO_{2max}$ .<sup>27,28</sup> The results of these studies showed that 12 months exercise at an intensity of 50–95% of  $VO_{2max}$  carried out three to five times per week produced an improvement of 37–42%. The longer training period, along with the large dispersion in intensity and various numbers of training sessions per week, makes these results difficult to compare to the present study. These studies however demonstrate that high aerobic exercise intensity is associated with a large improvement of  $VO_{2max}$ .

The aerobic exercise program was chosen to be aerobic interval exercise because this training method has been employed by our research

group in healthy individuals, yielding great improvements of  $VO_{2max}$  in a relative short time period.<sup>25</sup> This program was selected because it is typically used in training studies involving CAD patients.<sup>14</sup> Adachi et al,<sup>30</sup> compared 29 patients with previous myocardial infarction performing walking exercise over eight weeks. The  $VO_{2max}$  increased by 17%. These findings support our study with regard to the fact that aerobic exercise being more suitable for increasing  $VO_{2max}$  compared to higher intensity exercise. Contrary to these studies, Blumenthal et al<sup>31</sup> did not detect differences between moderate intensity and low intensity exercise after 12 weeks of training among 45 patients with myocardial infarction. The  $VO_{2max}$  increased by 11% within the high intensity group and 14% in the low intensity group, but the differences were not statistically significant. In sum, these studies indicate that aerobic exercise is more suitable for increasing  $VO_{2max}$ .

The initial  $VO_{2max}$  of the patients in our study was  $35.2 \pm 4.32$  ml/kg per min, which is higher compared to the other studies evaluating aerobic exercise in cardiac patients (18.7–25.3 ml/kg per min).<sup>29-31</sup> Thus, the improvement of  $VO_{2max}$  after considerable training when calculating percentage improvement from such a higher baseline value. In fact,  $VO_{2max}$  increased after training. To ensure that exercise is conducted at the proper intensity, it is important that patients are exercising close to their maximal effort on the initial  $VO_{2max}$ -test. If not, the reported exercise intensities are likely overestimates of the actual ranges. Hence, the term  $VO_{2max}$  was used instead of  $VO_{2peak}$  to describe exercise capacity throughout the study. The researchers state that use of the Bruce protocol with relatively large and uneven work increments, made several subjects exceeding their anaerobic threshold in the first 3-min stage.<sup>29</sup> Work rate increments that are too rapid may result in reduced exercise capacity and it is suggested that individualized protocols with estimated test duration of 8–12 minutes are optimal.<sup>24</sup> Adachi et al,<sup>30</sup> carried out the  $VO_{2max}$ -test on bicycle ergometers, which are reported to produce  $VO_{2max}$ -values that are

6–25% lower compared to treadmill exercise.<sup>24</sup> However, aerobic exercise is clearly beneficial in lowering mortality compared to a sedentary lifestyle, and current guidelines suggest that the incidence of sudden cardiac arrest across a variety of activities, except jogging, is similar to that expected by chance alone.<sup>9</sup> Exercise is also shown to be a potent trigger of myocardial infarction.<sup>33-35</sup> The adjusted relative risk of myocardial infarction during or soon after exertion have been found greater in persons who do not regularly participate in physical activity<sup>34,35</sup> and it is thus of great importance to get this group more active. However, selected exercise testing should be performed at the discretion of a physician before vigorous exercise in patients with known cardiovascular problems.<sup>36</sup> For stable CAD-patients in particular, Hauer et al,<sup>37,38</sup> demonstrated that adherence to prescribed target heart rate up to 95% of  $HR_{peak}$  ( $90\% VO_{2peak}$ ) reached during symptom-limited exercise testing is associated with very few ischaemic episodes even during high intensity exercise training. More studies are however needed to evaluate the comparative efficacy and safety of possible detrimental effects of high intensity exercise vs. other modes of exercise in a non-selected population of CAD-patients.

As increasing  $VO_{2max}$  is found to be a major determinant of increasing functional capacity and thereby survival,<sup>12,13</sup> this type of exercise may thus be employed to optimize the exercise component of rehabilitation programmes for stable CAD patients in the future.

## REFERENCES

1. Paffenbarger RS Jr, Hyde RT, Wing AL, Hsieh CC. Physical activity, all-cause mortality and longevity of college alumni. *N Engl J Med* 1986;314:605-13.
2. Oldridge NB, Guyatt GH, Fischer ME, Rimm AA. Cardiac rehabilitation after myocardial infarction. Combined experience of randomized clinical trials. *JAMA* 1988;260:945-50.
3. Blair SN, Kohl HW III, Paffenbarger RS Jr, Clark DG, Cooper KH, Gibbons LW. Physical fitness and all-cause mortality. A prospective study of healthy men and women. *JAMA* 1989;262:2395-401.

4. Farrell SW, Kampert JB, Kohl HW III, Barlow CE, Macera CA, Paffenbarger RS Jr, et al. Influences of cardiorespiratory fitness levels and other predictors on cardiovascular disease mortality in men. *Med Sci Sports Exerc* 1998;30:899-905.
5. O'Connor GT, Buring JE, Yusuf S, Goldhaber SZ, Olmstead EM, Paffenbarger RS, et al. An overview of randomized trials of rehabilitation with exercise after myocardial infarction. *Circulation* 1989;80:234-44.
6. Jolliffe JA, Rees K, Taylor RS, Thompson D, Oldridge N, Ebrahim S. Exercise-based rehabilitation for coronary heart disease. *Cochrane Database Syst Rev* 2001;1:CD001800.
7. American College of Sports Medicine Position Stand. Exercise for patients with coronary artery disease. *Med Sci Sports Exerc* 1994;26:i-v.
8. Fletcher GF, Balady G, Blair SN, Blumenthal J, Caspersen C, Chaitman B, et al. Statement on exercise: benefits and recommendations for physical activity programs for all Americans. A statement for health professionals by the Committee on Exercise and Cardiac Rehabilitation of the Council on Clinical Cardiology, American Heart Association. *Circulation* 1996;94:857-62.
9. Fletcher GF, Balady GJ, Amsterdam EA, Chaitman B, Eckel R, Fleg J, et al. Exercise standards for testing and training: a statement for healthcare professionals from the American Heart Association. *Circulation* 2001;104:1694-1740.
10. Shephard RJ, Balady GJ. Exercise as cardiovascular therapy. *Circulation* 1999;99:963-72.
11. Ades PA. Cardiac rehabilitation and secondary prevention of coronary heart disease. *N Engl J Med* 2001;345:892-902.
12. Myers J, Prakash M, Froelicher V, Do D, Partington S, Atwood JE, et al. Exercise capacity and mortality among men referred for exercise testing. *N Engl J Med* 2002;346:793-801.
13. Kavanagh T, Mertens DJ, Hamm LF, Beyene J, Kennedy J, Corey P, et al. Prediction of long-term prognosis in 12169 men referred for cardiac rehabilitation. *Circulation* 2002;106:666-71.
14. Swain DP, Franklin BA. Is there a threshold intensity for aerobic training in cardiac patients? *Med Sci Sports Exerc* 2002;34:1071-5.
15. Froelicher V, Jensen D, Genter F, Sullivan M, McKirnan MD, Witzum K, et al. A randomized trial of exercise training in patients with coronary heart disease. *JAMA* 1984;252:1291-7.
16. Hambrecht R, Niebauer J, Marburger C, Grunze M, Kalberer B, Hauer K, et al. Various intensities of leisure time physical activity in patients with coronary artery disease: effects on cardiorespiratory fitness and progression of coronary atherosclerotic lesions. *J Am Coll Cardiol* 1993;22:468-77.
17. Schuler G, Hambrecht R, Schlierf G, Niebauer J, Hauer K, Neumann J, et al. Regular physical exercise and low-fat diet. Effects on progression of coronary artery disease. *Circulation* 1992;86:1-11.
18. Haskell WL, Alderman EL, Fair JM, Maron DJ, Mackey SF, Superko HR, et al. Effects of intensive multiple risk factor reduction on coronary atherosclerosis and clinical cardiac events in men and women with coronary artery disease. The Stanford Coronary Risk Intervention Project (SCRIP). *Circulation* 1994;89:975-90.
19. Niebauer J, Hambrecht R, Velich T, Hauer K, Marburger C, Kalberer B, et al. Attenuated progression of coronary artery disease after 6 years of multifactorial risk intervention: role of physical exercise. *Circulation* 1997;96:2534-41.
20. Manson JE, Greenland P, LaCroix AZ, Stefanick ML, Mouton CP, Oberman A, et al. Walking compared with vigorous exercise for the prevention of cardiovascular events in women. *N Engl J Med* 2002;347:716-25.
21. Murphy M, Nevill A, Neville C, Biddle S, Hardman A. Accumulating brisk walking for fitness, cardiovascular risk, and psychological health. *Med Sci Sports Exerc* 2002;34:1468-74.
22. Tanasescu M, Leitzmann MF, Rimm EB, Willett WC, Stampfer MJ, Hu FB. Exercise type and intensity in relation to coronary heart disease in men. *JAMA* 2002;288:1994-2000.
23. Lee IM, Sesso HD, Oguma Y, Paffenbarger RS Jr. Relative intensity of physical activity and risk of coronary heart disease. *Circulation* 2003;107:1110-6.
24. Froelicher VF, Myers JN. Exercise and the heart. WB Saunders Company; 2000;11-38.
25. Helgerud J, Engen LC, Wisloff U, Hoff J. Aerobic endurance training improves soccer performance. *Med Sci Sports Exerc* 2001;33:1925-31.
26. Borg GA. Psychophysical bases of perceived exertion. *Med Sci Sports Exerc* 1982;14:377-81.
27. Ehsani AA, Martin WH III, Heath GW, Coyle EF. Cardiac effects of prolonged and intense exercise training in patients with coronary artery disease. *Am J Cardiol* 1982;50:246-54.
28. Ehsani AA, Biello DR, Schultz J, Sobel BE, Holloszy JO. Improvement of left ventricular contractile function by exercise training in patients with coronary artery disease. *Circulation* 1986;74:350-8.
29. Jensen BE, Fletcher BJ, Rupp JC, Fletcher GF, Lee JY, Oberman A. Training level comparison study. Effect of high and low intensity exercise on ventilatory threshold in men with coronary artery disease. *J Cardiopulm Rehabil* 1996;16:227-32.
30. Adachi H, Koike A, Obayashi T, Umezawa S, Aonuma K, Inada M. Does appropriate endurance exercise training improve cardiac function in patients with prior myocardial infarction? *Eur Heart J* 1996;17:1511-21.
31. Blumenthal JA, Rejeski WJ, Walsh-Riddle M, Emery CF, Miller H, Roark S, et al. Comparison of high- and low-intensity exercise training early after acute myocardial infarction. *Am J Cardiol* 1988;61:26-30.

32. Astrand PO, Rodahl K. Textbook of work physiology. Physiological bases of exercise. Third edition. New York: McGraw-Hill; 1986;295-353.
33. Tofler GH, Muller JE, Stone PH, Forman S, Solomon RE, Knatterud GL, et al. Modifiers of timing and possible triggers of acute myocardial infarction in the Thrombolysis in Myocardial Infarction Phase II (TIMI II) Study Group. *J Am Coll Cardiol* 1992;20:1049-55.
34. Mittleman MA, Maclure M, Tofler GH, Sherwood JB, Goldberg RJ, Muller JE. Triggering of acute myocardial infarction by heavy physical exertion. Protection against triggering by regular exertion. Determinants of Myocardial Infarction Onset Study Investigators. *N Engl J Med* 1993;329:1677-83.
35. Willich SN, Lewis M, Lowel H, Arntz HR, Schubert F, Schroder R. Physical exertion as a trigger of acute myocardial infarction. Triggers and Mechanisms of Myocardial Infarction Study Group. *N Engl J Med* 1993;329:1684-90.
36. Thompson PD, Buchner D, Pina IL, Balady GJ, Williams MA, Marcus BH, et al. Exercise and physical activity in the prevention and treatment of atherosclerotic cardiovascular disease: a statement from the Council on Clinical Cardiology (Subcommittee on Exercise, Rehabilitation, and Prevention) and the Council on Nutrition, Physical Activity, and Metabolism (Subcommittee on Physical Activity). *Circulation* 2003;107:3109-116.
37. Hauer K, Niebauer J, Weiss C, Marburger C, Hambrecht R, Schlierf G, et al. Myocardial ischemia during physical exercise in patients with stable coronary artery disease: predictability and prevention. *Int J Cardiol* 2000;75:179-86.
38. Wenger HA, Bell GJ. The interactions of intensity, frequency and duration of exercise training in altering cardiorespiratory fitness. *Sports Med* 1986;3:346-56.