



Evaluating the Neuroprotective Effect of Melatonin on Patients with Hemorrhagic Stroke Using Serum S100B Protein as a Prognostic Marker

Hamidreza Sharifnia¹, Mojtaba Mojtahedzadeh², Mehrnoush Dianatkah^{3,*}, Atabak Najafi¹, Arezoo Ahmadi¹, Farhad Najmeddin⁴, Minoos Dianatkah⁴, Nafiseh Alizadeh⁵, Atefeh Jafari⁶, Mandana Izadpanah⁷, Shahram Parvin⁸ and Maryam Daei⁹

¹Department of Anesthesiology and Critical Care, Sina Hospital, Tehran University of Medical Sciences, Tehran, Iran

²Department of Clinical Pharmacy, Tehran University of Medical Sciences, Tehran, Iran

³Department of Clinical Pharmacy, Isfahan University of Medical Sciences, Isfahan, Iran

⁴Heart Failure Research Center, Cardiovascular Research Institute, Isfahan University of Medical Sciences, Isfahan, Iran

⁵Baharloo Hospital, Tehran University of Medical Sciences, Tehran, Iran

⁶Department of Clinical Pharmacy, Rasht University of Medical Sciences, Rasht, Iran

⁷Department of Clinical Pharmacy, Ahvaz Jundishapur University of Medical Sciences, Ahvaz, Iran

⁸Chemical Injuries Research Center, Proteomics Laboratory, Baqiyatollah University of Medical Sciences, Tehran, Iran

⁹Department of Clinical Pharmacy, Alborz University of Medical Sciences, Alborz, Iran

*Corresponding author: Department of Clinical Pharmacy, Isfahan University of Medical Sciences, Isfahan, Iran. Email: mehrnoush.dianatkah@gmail.com

Received 2017 December 27; Revised 2018 December 29; Accepted 2019 January 23.

Abstract

Background: Intracerebral hemorrhage (ICH) is one of the most debilitating kinds of stroke. Recent evidence shows that the proper initiation of neuroprotective agents might save at risk neurons and improve the outcome.

Objectives: The focus of this study is to evaluate the neuroprotective effect of melatonin on patients with hemorrhagic stroke.

Methods: Forty adult patients with confirmed nontraumatic ICH, who were admitted to the ICU within 24 hours of the stroke onset were enrolled in this study. Subjects in the melatonin group received 30 mg of melatonin every night for 5 consecutive nights. In order to evaluate the intensity of the neuronal injury, S100B was assessed once on day 1 and, day 5 post ICU admission. Additionally, the length of ICU stay, mortality, and the duration of mechanical ventilation were also recorded.

Results: Forty patients completed the study. In both groups the plasma concentrations of S100B decreased after 5 days compared with their baseline values. However, this reduction was more significant in the melatonin compared to the control group (P-value < 0.05). The duration of mechanical ventilation and length of ICU stay was shorter in the melatonin group, and this difference was statistically significant for the length of ICU stay (P-value < 0.05), and marginally significant for the duration of mechanical ventilation (P-value = 0.065). The in-ICU mortality rate of the melatonin group was 15%, almost half of that of the control group (30%). However, this difference was not statistically significant.

Conclusions: In conclusion, melatonin can be considered as a harmless and effective neuroprotective agent with some unique features which has made it an appropriate adjunctive medicine for critically ill intubated patients.

Keywords: Melatonin, S100b, Hemorrhagic Stroke

1. Background

Spontaneous, nontraumatic intracerebral hemorrhage (ICH) is one of the most debilitating kinds of stroke (1, 2). Research has shown that most patients who are admitted with small ICHs, can survive if they receive appropriate medical care. Therefore, it can be concluded that good medical interventions may improve the outcome of these patients (3). Trials which have evaluated medical therapy for hemorrhagic stroke are very scarce, and the

optimal treatment strategy remains to be found.

Neuroprotective agents have not been extensively investigated in animal models of ICH. However, recent evidence shows that the proper initiation of neuroprotective agents might save at-risk neurons and improve outcomes in patients with hemorrhagic stroke (4). In recent years, melatonin has generated a great deal of interest as a neuroprotective agent for different neurological disorders (5) because of its low toxicity, antioxidative, antiapoptotic, and anti-inflammatory properties (5, 6). In contrast to the par-

tial distribution of other antioxidants, melatonin diffuses easily across the cell membranes and can act as an intracellular antioxidant as well as pass through the blood-brain barrier (7, 8). With regards to its extraordinary high antioxidant effect, numerous animal studies have been designed to evaluate the protective effect of melatonin on stroke (9).

The pathogenicity of stroke is a multifactorial process involving the interaction of multiple mechanisms, such as oxidative stress, excitotoxicity, mitochondrial dysfunction, and calcium dyshomeostasis (5). It has been widely accepted that oxidative stress and inflammation are the main causes of cell death in stroke (5). Because of its antioxidant effect, melatonin has shown to be protective against oxidative injury in various *in vivo* and *in vitro* models of neurodegenerative diseases (9-11). The protective effect of melatonin on animal models of stroke has been indicated in a metaanalysis (12).

In our research, S100B has been considered as a biomarker of damage associated molecular pattern (DAMP). This protein increases during the acute phase of brain damage (13). The serum levels of S100B have been shown to correlate with the severity of injury and outcome. Moreover, it is a useful tool for predicting the efficiency of the treatment and prognosis (14, 15). It has been reported that after a hemorrhagic stroke, plasma levels of S100B increase within 6 hours, peak at 24 hours, and then decline gradually over 2 days (15).

Previous studies have shown that pineal calcification and low melatonin levels are associated with symptomatic cerebral infarction and intracerebral hemorrhage (16, 17).

2. Objectives

The aim of this clinical trial was to evaluate the neuroprotective effect of exogenous melatonin on patients with hemorrhagic stroke.

3. Methods

Forty adult patients with confirmed acute ICH who were admitted to the ICU within 24 hours of stroke onset were enrolled in this randomized double-blind clinical trial.

Exclusion criteria were evidence of traumatic ICH, brain neoplasm, contraindications to receiving oral medication, Glasgow Coma scale (GCS) score of 8 or more, renal insufficiency (estimated glomerular filtration rate < 60 mL/min), and pregnancy or breastfeeding. In order to avoid the short-lasting pseudo increase in the plasma levels of S100B after surgery, a 24 hour interval between surgery and blood sampling was considered. If it was not

possible, then the patient would be excluded from the study (18, 19).

The written informed consent was signed by a legal surrogate instead of the patients due to the reduced level of consciousness. The study protocol was approved by the Ethics Committee for Human Research at Tehran University of Medical Sciences (code: IR.TUMS.VCR.REC.1395.884). Patients were randomized using the permuted-block randomization method. Patients in the treatment group received 30 mg of melatonin (Melatonin, Webber naturals, Canada) (as a tablet) every night throughout their nasogastric tube for 5 nights. Clinical characteristics of patients and demographic data were obtained at the time of enrollment. Additionally, mortality, the duration of mechanical ventilation, and the length of ICU stay were also recorded for all patients. The period of study was about one year.

S100B (using a Human S100B ELISA kit, Biovender), as a biomarker of neuronal injury was assessed once on the first day and day 5 post ICU admission. Blood samples (5 cc) were drawn from all patients through the antecubital vein into the sampling tube on days 1 and day 5. After waiting 10 minutes for coagulation, samples were centrifuged (10 min, 1000 rpm) in order to obtain serum, which was subsequently stored at -70°C until analyzed.

Student's *t* or Mann-Whitney U-test was used to compare the continuous variables between groups. Categorical variables were described by frequency in percentage, and the chi-square test was used to compare the groups. Analysis of covariance (ANCOVA) was used to compare the two groups when the baseline variable was assumed as a covariate. P-values less than or equal to 0.05 were considered significant. The IBM SPSS Statistics for Windows, version 20.0 (Armonk, NY: IBM Corp.) was used to conduct the analyses.

4. Results

Forty patients who met our inclusion criteria were enrolled in the study: twenty (50%) patients were allocated in the melatonin group and 20 (50%) patients in the control group (Figure 1). No significant difference was seen in the baseline characteristics of patients except for body mass index (BMI) (P-value = 0.049) (Table 1).

There was not any significant difference in the baseline plasma concentration of S100B between the two groups. In both groups the plasma concentrations of S100B were reduced after 5 days compared with their baseline values. However, this reduction was more significant in the melatonin compared to the control group (P-value < 0.05) (Table 2).

The length of ICU stay and the duration of mechanical ventilation was shorter in patients who received melatonin.




Figure 1. Consort flow chart of the study

tonin in comparison with the control group, and this difference was statistically significant for the length of ICU stay, and marginally significant for the duration of mechanical ventilation (P-value: 0.065) (Table 3).

A smaller proportion of the participants in the melatonin group were diagnosed with sepsis during their ICU stay in comparison with the control group. However, this difference was not statistically significant.

The in-ICU mortality rate of the melatonin group was 15%, almost half of that of the control group (30%). However, this difference was not statistically significant (Table 3).

5. Discussion

Melatonin has been studied in different fields of neurological disease such as seizures, brain edema, and traumatic brain injury with different dosing methods ranged from 3 mg to 50 mg, without any adverse effects (10).

The result of our research has shown that melatonin administered in a high dose within 24 hours of insult, can increase the clearance of S100B. Previous studies have

demonstrated that the therapeutic window for neuroprotection in humans is up to four days after stroke onset. With respect to this issue, an eligibility criteria of including patients within 24 hours of onset seems rational (4). Since it has been suggested that ischemic damage may continue for several days after infarction, 5 days of melatonin therapy has been considered in this clinical trial (4).

Available clinical data show that melatonin possesses hypnotic, anxiolytic, and analgesic properties, with minimal adverse effects on psychomotor performance, sleep wake cycle, and respiratory system. Moreover, it has been reported that melatonin is effective for both preventing and treating ICU delirium (20, 21).

The result of this randomized clinical trial revealed that administration of 30 mg of melatonin daily for 5 days is safe and effective in reducing S100B protein levels in patients with hemorrhagic stroke. One study has shown that in spite of critical illness, the bioavailability of melatonin after oral administration is satisfying, and the intestinal absorption is comparable to that of normal volunteers (22). The dose of 30 mg was chosen since investigations have shown that the antioxidant effect of melatonin

Table 1. Demographic and Clinical Characteristics of the Patients in Groups^{a, b}

	Melatonin (N = 20)	Control (N = 20)	P-Value
Age, y	57.7 ± 12.7	52.9 ± 13.7	0.329
Sex, male	12 (80)	9 (60)	0.232
BMI, kg/m ²	26.03 ± 1.84	24.6 ± 1.79	0.049 ^c
Hb, g/dL	12.1 ± 1.37	11.2 ± 1.52	0.245
Plt, 10 ³ /mm ³	195.0 ± 28.9	229.3 ± 35.9	0.061
Urea, mg/dL	33.9 ± 14.1	25.7 ± 9.27	0.215
Cr, mg/dL	0.96 ± 0.21	0.91 ± 0.15	0.634
INR	1.31 ± 0.25	1.26 ± 0.15	0.659
PTT, s	35.7 ± 9.05	38.2 ± 4.45	0.439
Ca, mg/dL	9.29 ± 0.85	8.94 ± 0.23	0.324
SOFA score	6.27 ± 0.70	6.64 ± 0.93	0.228
Apachi score	17.60 ± 4.22	16.9 ± 4.73	0.690
GCS day 1	6.13 ± 1.24	6.14 ± 0.95	0.982

Abbreviation: SOFA, The Sequential Organ Failure assessment.

^aValues are expressed as No. (%) or mean ± SD.

^bStudent's t-test and Chi-square were used to compare these values respectively.

^cP-value < 0.05 considered as significant.

Table 2. Effect of Melatonin on Serum Concentration of S100B^{a, b}

	Melatonin	Control	P-Value
S100-before, pg/mL	133.8 ± 33.5	132.6 ± 33.5	0.910
S100-after, pg/mL	58.7 ± 9.04	80.6 ± 20.3	< 0.001
Difference	52.0 ± 16.2	75.2 ± 28.5	0.003
P-value	< 0.001 ^c	< 0.001 ^c	< 0.001

^aValues are expressed as mean ± SD to compare values before and after intervention and between groups, paired t-test and t-test were used respectively.

^bANCOVA was used to compare values between groups when the baseline values were assumed as a covariate.

^cSignificant.

Table 3. Comparison of Duration of Mechanical Ventilation, Length of ICU Stay, in-ICU Mortality Rate, and Occurrence of Sepsis^a

	Melatonin	Control	P-Value
Duration of mechanical ventilation, d	4 (2-16)	12 (4-20)	0.065
Length of ICU stay, d	8 (6-21)	12 (8-25)	0.041
In ICU mortality	3 (15.0)	6 (30.0)	0.451
Sepsis	5 (25.0)	10 (50.0)	0.095
GCS day 5	8.60 ± 1.76	7.07 ± 1.33	0.064

^aNumerical values were reported as median (inter quartile range), and nominal factors as number (%), Mann-Whitney U-test and chi-square test were applied to compare these values respectively.

can be seen at doses above 10 mg, and with respect to safety, most of the studies usually use the dose as high as 20 mg to 50 mg (23).

Because there is strong evidence that oxidative stress is responsive to neuronal injury and its severity, antioxidant therapy may be extremely effective in reducing the cellular damage caused by free radicals (8). Moreover, it has been suggested that melatonin can exert some protection against neurotoxicity via other mechanisms, including regulation of calcium level, antiapoptotic activity, inhibition of mitochondrial permeability, decreasing cerebral edema, anticonvulsive, and anti-inflammatory effect (5, 11, 24, 25).

As has been shown in Table 2, a smaller proportion of the participants in the melatonin group were diagnosed with sepsis during their ICU stay in comparison with the control group. The protective effect of melatonin against sepsis and septic shock has been well documented in both human and animal models. It has been reported that melatonin can prevent circulatory failure, mitochondrial damage, and multi-organ failure, reduce lipid peroxidation and proinflammatory cytokines, and inhibit nitric oxide synthase (23, 26, 27). An investigation indicated that circadian rhythm was impaired in the septic patients when compared to the nonseptic ones, and it was concluded that reduced circadian melatonin secretion in septic patients may be the result of severe sepsis (28). Moreover, another theory is that reduced plasma melatonin concentrations may reflect the consumption of melatonin as an antioxidant (28, 29). In another study, decreased serum melatonin levels were associated with higher mortality rates in septic patients (30). A recent study revealed that the administration of melatonin as an adjuvant medicine in the treatment of septic newborns was associated with improvement of laboratory data and clinical outcomes (31).

As can be concluded from Table 2, the duration of mechanical ventilation was shorter in the melatonin group compared with the control one. Although this difference was not statistically significant, it can be hypothesized that melatonin may accelerate the weaning process by reducing the need for high doses of sedatives with respiratory depressant effects, inhibiting ventilator-associated lung injury, and improving the neurological status (32-34).

With regard to safety issues, experimental animal studies have claimed that melatonin is generally safe even in a dose as high as 200 mg/kg/day, and no serious adverse effects have been reported. Additionally, the safety of melatonin in humans has been shown in a metanalysis (35).

Our study did have some limitations. In this study, the total antioxidant capacity was not measured before and after the intervention in order to see to what extent the administration of the supplement has changed the total antioxidant capacity. Additionally, the absence of a placebo group in this study can adversely affect the results.

5.1. Conclusions

In conclusion, melatonin can be considered as a harmless and effective neuroprotective agent with some unique features which has made it an appropriate adjunctive medicine for critically ill intubated patients.

Footnotes

Authors' Contribution: None declared by the authors.

Clinical Trial Registration Code: This randomized clinical trial was registered in IRCT with an allocated number of IRCT2016100530164N1.

Conflict of Interests: We have no financial interests related to the material in the manuscript.

Ethical Approval: The ethical approval code was IR.TUMS.VCR.REC.1395.884.

Funding/Support: No funding and support.

References

- Sacco S, Marini C, Toni D, Olivieri L, Carolei A. Incidence and 10-year survival of intracerebral hemorrhage in a population-based registry. *Stroke*. 2009;**40**(2):394–9. doi: [10.1161/STROKEAHA.108.523209](https://doi.org/10.1161/STROKEAHA.108.523209). [PubMed: [19038914](https://pubmed.ncbi.nlm.nih.gov/19038914/)].
- Silva Y, Leira R, Tejada J, Lainez JM, Castillo J, Davalos A, et al. Molecular signatures of vascular injury are associated with early growth of intracerebral hemorrhage. *Stroke*. 2005;**36**(1):86–91. doi: [10.1161/01.STR.0000149615.51204.0b](https://doi.org/10.1161/01.STR.0000149615.51204.0b). [PubMed: [15550687](https://pubmed.ncbi.nlm.nih.gov/15550687/)].
- Hemphill J3, Greenberg SM, Anderson CS, Becker K, Bendok BR, Cushman M, et al. Guidelines for the Management of Spontaneous Intracerebral Hemorrhage: A Guideline for Healthcare Professionals From the American Heart Association/American Stroke Association. *Stroke*. 2015;**46**(7):2032–60. doi: [10.1161/STR.0000000000000069](https://doi.org/10.1161/STR.0000000000000069). [PubMed: [26022637](https://pubmed.ncbi.nlm.nih.gov/26022637/)].
- Dorman PJ, Sandercock PA. Considerations in the design of clinical trials of neuroprotective therapy in acute stroke. *Stroke*. 1996;**27**(9):1507–15. doi: [10.1161/01.STR.27.9.1507](https://doi.org/10.1161/01.STR.27.9.1507). [PubMed: [8784121](https://pubmed.ncbi.nlm.nih.gov/8784121/)].
- Andrabi SS, Parvez S, Tabassum H. Melatonin and Ischemic Stroke: Mechanistic Roles and Action. *Adv Pharmacol Sci*. 2015;**2015**:384750. doi: [10.1155/2015/384750](https://doi.org/10.1155/2015/384750). [PubMed: [26435711](https://pubmed.ncbi.nlm.nih.gov/26435711/)]. [PubMed Central: [PMC4575994](https://pubmed.ncbi.nlm.nih.gov/PMC4575994/)].
- Lekic T, Hartman R, Rojas H, Manaenko A, Chen W, Ayer R, et al. Protective effect of melatonin upon neuropathology, striatal function, and memory ability after intracerebral hemorrhage in rats. *J Neurotrauma*. 2010;**27**(3):627–37. doi: [10.1089/neu.2009.1163](https://doi.org/10.1089/neu.2009.1163). [PubMed: [20350200](https://pubmed.ncbi.nlm.nih.gov/20350200/)]. [PubMed Central: [PMC2867555](https://pubmed.ncbi.nlm.nih.gov/PMC2867555/)].
- Dominguez-Rodriguez A, Abreu-Gonzalez P, Garcia-Gonzalez MJ, Kaski JC, Reiter RJ, Jimenez-Sosa A. A unicenter, randomized, double-blind, parallel-group, placebo-controlled study of Melatonin as an Adjunct in patients with acute myocardial Infarction undergoing primary Angioplasty The Melatonin Adjunct in the acute myocardial Infarction treated with Angioplasty (MARIA) trial: study design and rationale. *Contemp Clin Trials*. 2007;**28**(4):532–9. doi: [10.1016/j.cct.2006.10.007](https://doi.org/10.1016/j.cct.2006.10.007). [PubMed: [17123867](https://pubmed.ncbi.nlm.nih.gov/17123867/)].
- Gonzales-Portillo GS, Lozano D, Aguirre D, Reyes S, Borlongan CV, Tajiri N, et al. An update on the use of melatonin as a stroke therapeutic. *Minerva Med*. 2015;**106**(3):169–75. [PubMed: [25000217](https://pubmed.ncbi.nlm.nih.gov/25000217/)].
- Watson N, Diamandis T, Gonzales-Portillo C, Reyes S, Borlongan CV. Melatonin as an Antioxidant for Stroke Neuroprotection. *Cell Transplant*. 2016;**25**(5):883–91. doi: [10.3727/096368915X689749](https://doi.org/10.3727/096368915X689749). [PubMed: [26497887](https://pubmed.ncbi.nlm.nih.gov/26497887/)].
- Altun A, Ugur-Altun B. Melatonin: therapeutic and clinical utilization. *Int J Clin Pract*. 2007;**61**(5):835–45. doi: [10.1111/j.1742-1241.2006.01191.x](https://doi.org/10.1111/j.1742-1241.2006.01191.x). [PubMed: [17298593](https://pubmed.ncbi.nlm.nih.gov/17298593/)].
- Hardeland R, Cardinali DP, Brown GM, Pandi-Perumal SR. Melatonin and brain inflammaging. *Prog Neurobiol*. 2015;**127**:46–63. doi: [10.1016/j.pneurobio.2015.02.001](https://doi.org/10.1016/j.pneurobio.2015.02.001). [PubMed: [25697044](https://pubmed.ncbi.nlm.nih.gov/25697044/)].
- Macleod MR, O'Collins T, Horky LL, Howells DW, Donnan GA. Systematic review and meta-analysis of the efficacy of melatonin in experimental stroke. *J Pineal Res*. 2005;**38**(1):35–41. doi: [10.1111/j.1600-079X.2004.00172.x](https://doi.org/10.1111/j.1600-079X.2004.00172.x). [PubMed: [15617535](https://pubmed.ncbi.nlm.nih.gov/15617535/)].
- Brunswick AS, Hwang BY, Appelboom G, Hwang RY, Piazza MA, Connolly EJ. Serum biomarkers of spontaneous intracerebral hemorrhage induced secondary brain injury. *J Neurol Sci*. 2012;**321**(1-2):1–10. doi: [10.1016/j.jns.2012.06.008](https://doi.org/10.1016/j.jns.2012.06.008). [PubMed: [22857988](https://pubmed.ncbi.nlm.nih.gov/22857988/)].
- Sanchez-Pena P, Pereira AR, Sourour NA, Biondi A, Lejean L, Colonne C, et al. S100B as an additional prognostic marker in subarachnoid aneurysmal hemorrhage. *Crit Care Med*. 2008;**36**(8):2267–73. doi: [10.1097/CCM.0b013e3181809750](https://doi.org/10.1097/CCM.0b013e3181809750). [PubMed: [18596638](https://pubmed.ncbi.nlm.nih.gov/18596638/)].
- Hu YY, Dong XQ, Yu WH, Zhang ZY. Change in plasma S100B level after acute spontaneous basal ganglia hemorrhage. *Shock*. 2010;**33**(2):134–40. doi: [10.1097/SHK.0b013e3181ad5c88](https://doi.org/10.1097/SHK.0b013e3181ad5c88). [PubMed: [19487970](https://pubmed.ncbi.nlm.nih.gov/19487970/)].
- Kitkhandee A, Sawanyawisuth K, Johns NP, Kanpittaya J, Johns J. Pineal calcification is associated with symptomatic cerebral infarction. *J Stroke Cerebrovasc Dis*. 2014;**23**(2):249–53. doi: [10.1016/j.jstrokecerebrovasdis.2013.01.009](https://doi.org/10.1016/j.jstrokecerebrovasdis.2013.01.009). [PubMed: [23434443](https://pubmed.ncbi.nlm.nih.gov/23434443/)].
- Kitkhandee A, Sawanyawisuth K, Johns J, Kanpittaya J, Tuntapakul S, Johns NP. Pineal calcification is a novel risk factor for symptomatic intracerebral hemorrhage. *Clin Neurol Neurosurg*. 2014;**121**:51–4. doi: [10.1016/j.clineuro.2014.03.019](https://doi.org/10.1016/j.clineuro.2014.03.019). [PubMed: [24793475](https://pubmed.ncbi.nlm.nih.gov/24793475/)].
- Uncu M. S100b protein levels in acute ischemic events. *Turk J Geriatr*. 2013;**16**(3):356–7.
- Schulte S, Podlog LW, Hamson-Utley JJ, Strathmann FG, Struder HK. A systematic review of the biomarker S100B: implications for sport-related concussion management. *J Athl Train*. 2014;**49**(6):830–50. doi: [10.4085/1062-6050-49.3.33](https://doi.org/10.4085/1062-6050-49.3.33). [PubMed: [25299445](https://pubmed.ncbi.nlm.nih.gov/25299445/)]. [PubMed Central: [PMC4264656](https://pubmed.ncbi.nlm.nih.gov/PMC4264656/)].
- Dianatkhah M, Ghaeli P, Hajhossein Talasaz A, Karimi A, Salehiomran A, Bina P, et al. Evaluating the Potential Effect of Melatonin on the post-Cardiac Surgery Sleep Disorder. *J Tehran Heart Cent*. 2015;**10**(3):122–8. [PubMed: [26697084](https://pubmed.ncbi.nlm.nih.gov/26697084/)]. [PubMed Central: [PMC4685367](https://pubmed.ncbi.nlm.nih.gov/PMC4685367/)].
- Maitra S, Baidya DK, Khanna P. Melatonin in perioperative medicine: Current perspective. *Saudi J Anaesth*. 2013;**7**(3):315–21. doi: [10.4103/1658-354X.115316](https://doi.org/10.4103/1658-354X.115316). [PubMed: [24015137](https://pubmed.ncbi.nlm.nih.gov/24015137/)]. [PubMed Central: [PMC3757807](https://pubmed.ncbi.nlm.nih.gov/PMC3757807/)].
- Gin W, Shaw RJ, Kay AB. Airways reversibility after prednisolone therapy in chronic asthma is associated with alterations in leukocyte function. *Am Rev Respir Dis*. 1985;**132**(6):1199–203. doi: [10.1164/arrd.1985.132.6.1199](https://doi.org/10.1164/arrd.1985.132.6.1199). [PubMed: [2934011](https://pubmed.ncbi.nlm.nih.gov/2934011/)].
- Kurdi MS, Patel T. The role of melatonin in anaesthesia and critical care. *Indian J Anaesth*. 2013;**57**(2):137–44. doi: [10.4103/0019-5049.111837](https://doi.org/10.4103/0019-5049.111837). [PubMed: [23825812](https://pubmed.ncbi.nlm.nih.gov/23825812/)]. [PubMed Central: [PMC3696260](https://pubmed.ncbi.nlm.nih.gov/PMC3696260/)].
- Rathnasamy G, Ling EA, Kaur C. Therapeutic implications of melatonin in cerebral edema. *Histol Histopathol*. 2014;**29**(12):1525–38. doi: [10.14670/HH-29.1525](https://doi.org/10.14670/HH-29.1525). [PubMed: [24876075](https://pubmed.ncbi.nlm.nih.gov/24876075/)].
- Arushanian EB. [Experimental and clinical evidence for anticonvulsant activity of melatonin]. *Zh Nevrol Psikhiatr Im S S Korsakova*. 2013;**113**(11):11–5. [PubMed: [24479166](https://pubmed.ncbi.nlm.nih.gov/24479166/)].
- Srinivasan V, Mohamed M, Kato H. Melatonin in bacterial and viral infections with focus on sepsis: a review. *Recent Pat Endocr Metab Immune Drug Discov*. 2012;**6**(1):30–9. doi: [10.2174/187221412799015317](https://doi.org/10.2174/187221412799015317). [PubMed: [22264213](https://pubmed.ncbi.nlm.nih.gov/22264213/)].

27. Galley HF, Lowes DA, Allen L, Cameron G, Aucott LS, Webster NR. Melatonin as a potential therapy for sepsis: a phase I dose escalation study and an ex vivo whole blood model under conditions of sepsis. *J Pineal Res.* 2014;**56**(4):427-38. doi: [10.1111/jpi.12134](https://doi.org/10.1111/jpi.12134). [PubMed: [24650045](https://pubmed.ncbi.nlm.nih.gov/24650045/)]. [PubMed Central: [PMC4279949](https://pubmed.ncbi.nlm.nih.gov/PMC4279949/)].
28. Mundigler G, Delle-Karth G, Koreny M, Zehetgruber M, Steindl-Munda P, Marktl W, et al. Impaired circadian rhythm of melatonin secretion in sedated critically ill patients with severe sepsis. *Crit Care Med.* 2002;**30**(3):536-40. doi: [10.1097/00003246-200203000-00007](https://doi.org/10.1097/00003246-200203000-00007). [PubMed: [11990911](https://pubmed.ncbi.nlm.nih.gov/11990911/)].
29. Bourne RS, Mills GH. Melatonin: possible implications for the postoperative and critically ill patient. *Intensive Care Med.* 2006;**32**(3):371-9. doi: [10.1007/s00134-005-0061-x](https://doi.org/10.1007/s00134-005-0061-x). [PubMed: [16477412](https://pubmed.ncbi.nlm.nih.gov/16477412/)].
30. Lorente L, Martin MM, Abreu-Gonzalez P, de la Cruz T, Ferreres J, Sole-Violan J, et al. Serum melatonin levels are associated with mortality in severe septic patients. *J Crit Care.* 2015;**30**(4):860 e1-6. doi: [10.1016/j.jcrc.2015.03.023](https://doi.org/10.1016/j.jcrc.2015.03.023). [PubMed: [25869726](https://pubmed.ncbi.nlm.nih.gov/25869726/)].
31. El Fragy M, El-Sharkawy HM, Attia GF. Use of melatonin as an adjuvant therapy in neonatal sepsis. *J Neonatal Perinatal Med.* 2015;**8**(3):227-32. doi: [10.3233/NPM-15814072](https://doi.org/10.3233/NPM-15814072). [PubMed: [26485549](https://pubmed.ncbi.nlm.nih.gov/26485549/)].
32. Gitto E, Reiter RJ, Sabatino G, Buonocore G, Romeo C, Gitto P, et al. Correlation among cytokines, bronchopulmonary dysplasia and modality of ventilation in preterm newborns: improvement with melatonin treatment. *J Pineal Res.* 2005;**39**(3):287-93. doi: [10.1111/j.1600-079X.2005.00251.x](https://doi.org/10.1111/j.1600-079X.2005.00251.x). [PubMed: [16150110](https://pubmed.ncbi.nlm.nih.gov/16150110/)].
33. Dessap AM, Roche-Campo F, Launay JM, Charles-Nelson A, Katsahian S, Brun-Buisson C, et al. Delirium and Circadian Rhythm of Melatonin During Weaning From Mechanical Ventilation: An Ancillary Study of a Weaning Trial. *Chest.* 2015;**148**(5):1231-41. doi: [10.1378/chest.15-0525](https://doi.org/10.1378/chest.15-0525). [PubMed: [26158245](https://pubmed.ncbi.nlm.nih.gov/26158245/)].
34. Mistraretti G, Umbrello M, Sabbatini G, Miori S, Taverna M, Cerri B, et al. Melatonin reduces the need for sedation in ICU patients: a randomized controlled trial. *Minerva Anesthesiol.* 2015;**81**(12):1298-310. [PubMed: [25969139](https://pubmed.ncbi.nlm.nih.gov/25969139/)].
35. Ramos E, Patino P, Reiter RJ, Gil-Martin E, Marco-Contelles J, Parada E, et al. Ischemic brain injury: New insights on the protective role of melatonin. *Free Radic Biol Med.* 2017;**104**:32-53. doi: [10.1016/j.freeradbiomed.2017.01.005](https://doi.org/10.1016/j.freeradbiomed.2017.01.005). [PubMed: [28065781](https://pubmed.ncbi.nlm.nih.gov/28065781/)].